

4TH INTERNATIONAL CONFERENCE:
**CONSTRUCTIVE
MATHEMATICAL
ANALYSIS**

ABSTRACT BOOK

01-02 JULY 2026

The Fourth International Conference: Constructive Mathematical Analysis

Abstract Book

July 01-02, 2026
KONYA, TÜRKİYE



**This conference is supported by the Scientific and Technological Research Council of
Türkiye (TÜBİTAK) 1001-Project 123F123.*

CMA
CONSTRUCTIVE MATHEMATICAL ANALYSIS

FOREWORDS

Dear Participants,

It's my pleasure to chair The International Conference on "4th International Conference: Constructive Mathematical Analysis". 4th International Conference: Constructive Mathematical Analysis is an activity of the journal Constructive Mathematical Analysis. The first two series of the event were organized as workshops. Based on the requests from international researchers studying on constructive mathematical analysis, this series was organized as a conference. Our 2025 conference was successfully completed, and this 2026 edition is organized with the valuable support and partnership of Selçuk University (Türkiye), Gazi University (Türkiye), Lamberto Cesari Inter. Res. Center (Italy), Vasyl Stefanyk Carpathian National University (Ukraine), Thammasat University (Thailand), ANA&A (Italy), and The Scientific and Technological Research Council of Türkiye (TÜBİTAK). Specifically, this conference is supported by the Scientific and Technological Research Council of Türkiye (TÜBİTAK) 1001-Project 123F123.

This year, the conference proudly features a "**Special Session Dedicated to the 75th Birthday of Prof. Ioan Raşa**". This session serves as a tribute to Prof. Raşa's outstanding achievements, invaluable contributions to mathematical science, and his inspiring academic legacy.

The main goal of this conference is to promote, encourage, and provide a forum for the academic exchange of ideas and recent research works on any field of Analysis and Function Theory. The conference presented new results and future challenges, in a series of keynote lectures and contributed short talks. We thank invited speakers distinguished **Prof. Carlo Bardaro**, distinguished **Prof. Raul Curto** and distinguished **Prof. Feng Dai** for contribution to the our conference.

The full texts of the papers presented in this conference will be published in *Springer Proceedings in Mathematics & Statistics*. The papers presented in this conference will be considered in the special issues of journals *ALTAY Conference Proceedings in Mathematics*, *Applied Mathematics*, *Carpathian Mathematical Publications*, *Constructive Mathematical Analysis*, *Gazi University Journal of Science*, *Mathematical and Computational Applications*, *Journal of Mathematical Sciences* and *Symmetry*.

This booklet contains the titles and abstracts of all invited and contributed talks at the *The Fourth International Conference: Constructive Mathematical Analysis* and is available on the conference website.

Prof. Dr. Tuncer ACAR
On Behalf of Organizing Committee - Chairman



*We warmly congratulate Prof. Ioan Raşa on his 75th birthday
and wish him a healthy, happy and long life.*

Chairman of Conference

Tuncer Acar
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
tunceracar@ymail.com

Conference Co-Chairs

Sadettin Kursun
Department of Information Security Technology, Faculty of Humanities and Social Sciences,
İzmir Kâtip Çelebi University, Türkiye
sadettinkursun@yahoo.com

Metin Turgay
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
metinturgay@yahoo.com

Organizing Committee

Özlem Acar
Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
acarozlem@ymail.com

Ali Aral
Kırıkkale University, Türkiye
aliaral73@yahoo.com

Sorina Barza
Karlstad University, Sweden
sorina.barza@kau.s

Michele Campiti
University of Salento, Italy
michele.campiti@unisalento.it

Danilo Costarelli
University of Perugia, Italy
danilo.costarelli@unipg.it

Roman Dmytryshyn
Vasyl Stefanyk Carpathian National University, Ukraine
dmytryshynr@hotmail.com

Borislav Radkov Draganov
Sofia University, Bulgaria
bdraganov@fmi.uni-sofia.bg

Gurhan Icoz
Gazi University, Türkiye
gurhanicoz@gazi.edu.tr

Mohamed A. Khamsi
University of Texas at El Paso, USA
mkhamsi@kfupm.edu.sa

Hendrik Van Maldeghem
University of Gent, Belgium
Hendrik.VanMaldeghem@UGent.be

Peter Massopust
Technical University of Munich, Germany
massopust@ma.tum.de

Donal O'Regan
National University of Ireland, Ireland
donal.oregan@nuigalway.ie

Wutiphol Sintunavarat
Thammasat University, Thailand
wutiphol@mathstat.sci.tu.ac.th

Calogero Vetro
Department of Mathematics, Faculty of Science, University of Palermo, Italy
calogero.vetro@unipa.it

Ferenc Weisz
Eötvös University, Hungary
weiszf@inf.elte.hu

Kehe Zhu
Department of Mathematics and Statistics State University of New York, USA
kzhu@math.albany.edu

Scientific Committee

Ulrich Abel
Mittelhessen University of Applied Sciences, Germany

Francesco Altomare
University of Bari Aldo Moro, Italy

Bayram Cekim
Gazi University, Türkiye

Clemente Cesarano
International Telematic University Uninettuno, Italy

Mirella Cappaletti Montano
University of Bari Aldo Moro, Italy

Feng Dai
University of Alberta, Canada

Harun Karşlı
Bolu Abant İzzet Baysal University, Türkiye

Fuad Kittenah
University of Jordan, Jordan

Vita Leonessa
University of Basilicata, Italy

Guozhen Lu
University of Connecticut, USA

Ilaria Mantellini
University of Perugia, Italy

Stefano de Marchi
University of Padova, Italy

Francisco J. Marcellan
Universidad Carlos III de Madrid, Spain

Manuel Lopez-Pellicer
University of Valencia, Spain

Ioan Raşa
Technical University of Cluj-Napoca, Romania

Salvador Romaguera Bonilla
Polytechnic University of Valencia, Spain

Anna Rita Sambucini
University of Perugia, Italy

Gianluca Vinti
University of Perugia, Italy

ICCMA

Secretary

Emre Bayrakcı

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
emre42bayrakci@gmail.com

Samet Değle

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
smt.degle@gmail.com

Tugce Delen

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
tugcedelen@yahoo.com

Dilek Erdogan

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
dilekozer@yahoo.com

Esref Erdogan

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
esreferdogan@yahoo.com

Serife Nur Karaman

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
serifenurkaraman@yahoo.com

Aybala Sevede Özkapu

Department of Mathematics, Faculty of Science, Selçuk University, Türkiye
aybalasevdeozkapu@yahoo.com

ICCMA

Contents

Exponential Sampling Theory: Error Estimates via Mellin Distance CARLO BARDARO	1
Classes of Operators Related to Subnormal Operators RAUL E. CURTO	2
Sampling Discretization of Integral Norms in Finite-Dimensional Spaces FENG DAI	3
Investigation of Physics-Informed Neural Networks for Solving the Allen-Cahn Equation ADIL NAURYZBAY, ZHANARS ABDIRAMANOV, ESIMKHAN ZHUMAZHANOV, DASTAN URNEK AND KAZEZ ERTAI	4
Modified Process Capability Index under Unit-Gumbel Distribution: Comparative Analysis of Point Estimators YUNUS AKDOGAN, TENZILE ERBAYRAM	5
Geometric Analysis of Magnetic Trajectories of Charged Particles and Their Relations with Associated Curves HÜDA NUR AKGÜL, MUHAMMED T. SARIAYDIN	6
OMNIAI: AI-Powered Chatbot-Based Appointment Management System MUHAMMET İKBAL AKGÜL, GÖKHAN UÇKAN AND EMRE ÇOMAK	7
On Integral Curves Generated by PAF in Euclidean 3-Space SELEN AKPINAR, MUHAMMED T. SARIAYDIN	8
Fixed Point Approximation Schemes for Wasserstein Gradient Flows in Generative Modeling OSMAN ALAGÖZ, EMRAH HASPOLAT	9
Two Dimensional Fuzzy Differential Transform Method for Solving of Two Dimensional Nonlinear Fuzzy Volterra Integral Equations ARTAN F. ALIDEMA, FREDERIK DARA	10
On the Approximation of Borel Derivatives by Nonlinear Szász Mirakyan Durrmeyer Operators HÜSEYİN ERHAN ALTIN	11
Almost Automorphic and Pseudo-Almost Automorphic Solutions: A Unified Theory with Applications to Classical and Fractional Evolution Systems DHAOU LASSOUED, JEHAD ALZABUT	12
Numerical Study on a Class of Higher-Order Singular Polytropic Equations DJIHENE ARAFA, NOURIA ARAR	13
On a New Modification of the Mellin-Gauss-Weierstrass Operators in the Weighted Mellin-Lebesgue Spaces ALI ARAL	14

On Some Matrix Transformations of Sequences of Function	15
ALI ARPACIOĞLU, TUNCAY TUNÇ	
A Neurodynamical Neural Network Approach for Solving Support Vector Machine Problems with Generalized Pinball Loss Functions via Hyperbolic Tangent Smoothing Technique	16
RATTANAPORN WANGKEEREE, RABIAN WANGKEEREE, YIRGA ABEBE BELAY AND KASAMSUK UNGCHITTRAKOOL	
Where/Why Do We Use the Bivariate/Multivariate Positive Linear/Nonlinear Operators?	17
MURAT BODUR	
Existence of Solutions for a Fractional $p(x, \cdot)$ -Laplacian Problem with Neumann Boundary Conditions on an Exterior Domain	18
ABDELKRIM BARBARA, AHMED BOUSMAHA AND MOHAMMED SHIMI	
Magic Squares: From Ancient Aesthetics to a New Recursive Construction Method	19
SYRGAK KYDYRALIEV, ANARKUL URDALETOVA, ELENA BUROVA AND SALTANAT JAPAROVA	
Eigenvalue Expansions for a Conformable Sturm–Liouville Operator with Mixed Boundary Conditions	20
OLGUN CABRI	
Appell-Vietoris Polynomials and Their Zeros	21
ISABEL CAÇÃO, MARIA IRENE FALCÃO, HELMUTH ROBERT MALONEK AND GRAÇA TOMAZ	
Approximation Processes on Minimal Surfaces	22
MICHELE CAMPITI	
Statistical Inference of a New Four-Parameter Marshall Olkin Distribution: Comprehensive Simulation and Measurement Applications	23
ERDEM CANKUT, KADIR KARAKAYA	
Inference for the Reliability and Mean Time to Failure of Protected Consecutive k -out-of- n :G Systems	24
ERDEM CANKUT, YUNUS AKDOĞAN, KADIR KARAKAYA AND COŞKUN KUŞ	
A Study on Associated Curves and Their Curvature Relations	25
ALEYNA ÇETINKAYA, MUHAMMED T. SARIAYDIN	
Stability to Perturbations of Modified S -Fractions with Applications	26
VOLODYMYR HLADUN, MARTA DMYTRYSHYN AND OLEKSANDR CHABAN	
A Chemotaxis-Navier-Stokes System with Dynamical Boundary Conditions	27
BAILI CHEN	
Korovkin-Type Approximation for Non-positive Operators	28

VIJAYALAKSHMI BALAKRISHNAN KIRAN KUMAR, MEMANA NEELAKANTAN NARAYANAN NAMBOODIRI AND VINAYA POYIL CHITTARICKAL	
A Generalized Fourier-Feynman Transform over Paths in Abstract Wiener Space DONG HYUN CHO	29
A Neurodynamic Optimization Model for Smooth Pinball Loss Based Support Vector Machines VIPAVEE DAMMINSED, PHAM THI VUI AND RABIAN WANGKEEREE	30
Automatic Differentiation Using Neural Networks and Their Comparison with Higher Order Spectral Differentiation MRINAL KANTI DAS, LOKENDRA KUMAR BALYAN	31
Fixed-Point Iterations for Analyzing the Topology of Biological Networks TUGCE DELEN, OZLEM ACAR	32
New Results on Fixed Point Theory in Perturbed Metric Spaces with Directed Graphs TUGCE DELEN, OZLEM ACAR	33
On the Approximations of the Ratios of Double Hypergeometric Functions by Branched Continued Fractions ROMAN DMYTRYSHYN	34
Solution Structure of Higher-Order Nonlinear Hadamard-Type Fractional p -Laplacian Problems on Finite Domains MÜRÜVVET TUVA ERBAŞ, ERBİL ÇETİN	35
XBilal Distribution: Properties, Parameter Estimation and Applications TENZILE ERBAYRAM, YUNUS AKDOĞAN	36
On a Statistical Versions of Certain Types of Alpha Convergence ALPER ERDEM, TUNCAY TUNÇ	37
Convergence and Asymptotic Behavior of Mellin-Steklov Exponential Sampling Series in Logarithmically Weighted Spaces DILEK ERDOĞAN, TUNCER ACAR	38
Approximation of Discontinuous Functions By Kantorovich Exponential Neural Network Op- erators DILEK ERDOĞAN, SERIFE NUR KARAMAN	39
Fixed Point Results for Rational Type F -Contractions in Perturbed Metric Spaces ESREF ERDOĞAN, OZLEM ACAR	40
Fixed Point Theorems for Z -Contractions in Perturbed Metric Spaces EŞREF ERDOĞAN, AYBALA SEVDE ÖZKAPU	41
An Affine ARX-Based Adaptive RST Framework for Disturbed Buck Converter Systems FATİH EVRAN	42

Dominants and Their Properties in Lattice Scaled Spaces NIYAZI ANIL GEZER	44
Numerical Stability of Branched Continued Fraction Expansions of Lauricella-Saran Hypergeometric Functions F_K and Their Ratios in Special Cases VITALIY GORAN	45
Some Properties of Generalized Degenerate Tangent Polynomials and Their Associated Matrices JUAN HERNÁNDEZ, YAMILET QUINTANA AND WILLIAM RAMÍREZ	46
Structural Stability and Rational q -Deformations in Parikh q -Determinants SEYDA ILDAN, AYNUR YALCINER	47
On the Zero Behavior of Degenerate Bernstein Basis Functions RODRIGO INGELMO, YAMILET QUINTANA	48
Approximation Properties of Modified Neural Network Operators Activated by Sigmoidal Functions SERIFE NUR KARAMAN, TUNCER ACAR	49
Convergence of Multidimensional Mellin Type Singular Integral Operators with Non-Isotropic Kernels HARUN KARS LI	50
Mid p -Summable Sequences and Applications to Lipschitz Operators AYŞEGÜL KETEN ÇOPUR	51
Numerical Spectral Analysis of a Chafee–Infante Type Equation on Unbounded Domains HADIL KHENISSA, NOURIA ARAR	52
On Exponential-Type Sampling Series: Upper Bounds of Operator Differences SADETTIN KURSUN	53
A Mixture Model Using Unit Type Lindley Distributions COŞKUN KUŞ, SÜMEYRA SERT	54
Convergence Domains of Branched Continued Fraction Expansions of Ratio $H_4(a, d + 1; c, d; \mathbf{z})/H_4(a + 1, d + 1; c, d + 1; \mathbf{z})$ ILONA-ANNA LUTSIV	55
Pseudospectra in Banach Algebras and Banach Jordan Algebras ABDELAZIZ MAOUCHE	56
Korovkin and Voronovskaya Results for Positive Bivariate Operators under Generalized Convergence FRANCISCO-JAVIER MARTÍNEZ-SÁNCHEZ, PEDRO GARRANCHO AND DANIEL CÁRDENAS-MORALES	57

Bridging Extending and Supplemented Modules: Towards a Unified Framework in Module Theory	58
ZAHRA MOAED ABD AL-MAJEED, ZAINAB RAAD SHAKER, TAMARA WATHIB MOHAMMED AND SUR A KHALEEL JAWAD	
A θ -Scheme for Complex Nonlinear Klein-Gordon Equations in the Nonrelativistic Limit: Stability Analysis and Numerical Results	59
AREZOO LATIF ALTOJAR, REZA MOKHTARI	
Convergence and Smoothness of the Mehler-Fock Transform and Application to Overdetermined Boundary Value Problems in $\mathbb{H}^n \times \mathbb{R}$	60
FILIPPO MORABITO	
A Constructive Definition of the Improper Riemann Integral on a Separable Banach Space	61
TIMOTHY MYERS	
On the Representation of Lauricella-Saran Hypergeometric Functions F_M and Their Ratios in Special Cases by Branched Continued Fractions	62
IVAN NYZHNYK	
A Randomized Shifted Finite Volume Method to Break Resonance Locking in 1D Homogenization	63
BIENVENU ONDAMI	
Fixed Point Theorems in Digital Metric Spaces and Applications in Fractal Image Compression	64
AYBALA SEVDE ÖZKAPU, ÖZLEM ACAR	
Approximation by a Novel Modification of a Mellin-Type Kernel	65
FIRAT OZSARAC	
Hyperbolic Wavelets, Basis Problem	66
MARGIT PAP	
A Smooth Lagrangian-Based Neural Network for Solving Semidefinite Programming	67
PAKKAPON PREECHASILP, RATTANAPORN WANGKEEREE, RABIAN WANGKEEREE, KASAMSUK UNGCHITTRAKOOL, YIRGA ABEBE BELAY AND PANATDA BOONMAN	
Rational Approximation and Meromorphic Continuation	68
VASILY PROKHOROV	
Degenerate Appell Sequences via Caputo-Type Fractional Difference Operator	69
STIVEN DÍAZ, WILLIAM RAMÍREZ, YAMILET QUINTANA AND CLEMENTE CESARANO	
Inequalities for Complex Polynomials	70
POONAM BABANI RANE, PRASANNA KUMAR	

Spherical Number Algebras: An Invariant-First Framework from Conceptual Architecture to Symbolic and Numeric Presentations	71
MOHAMMAD REZA ALAEI JORDEHI	
Sums as Sets	72
ANTHONY J. RIPA	
On the Oscillatory Behavior of Cesàro Means for Fourier-Jacobi Series	73
SAMRA SADIKOVIĆ HAJRIĆ, JASMINA MUMINOVIĆ HUREMOVIĆ	
Modeling Bounded Data for the Unit Transformed Inverse Weibull Distribution: Estimation and Application	74
SULE SAGLAM, KADIR KARAKAYA	
Reliability Evaluation for the Linear or Circular Consecutive k-out-of-n: G System Using Continuous Time Bayesian Network	75
RANGANATHAN SAKTHIVEL, GUNDALA VIJAYALAKSHMI AND SAHADEVAN SAK- THIPRIYA	
Modern Iterative Optimization with Monotone Operators and Nonexpansive Mappings	76
BERAT SERT, ÖZLEM ACAR	
A New Nonparametric k -Sample Test Based on Interval Counts of Order Statistics	77
SÜMEYRA SERT, COŞKUN KUŞ	
Fixed Point Theorems for Nonlinear Contractive Mappings in Cone-Valued θ -Type Multiplicative Metric Spaces	78
PRAVIN SINGH, VIRATH SINGH, SHIVANI SINGH AND SANI SALISU	
Dynamic Aspects of Composition Operators on Weighted Holomorphic Function Spaces	79
HAYRI TOPAL	
Simultaneous Approximation by Bivariate Kantorovich-Type Operators in Weighted Spaces	80
METIN TURGAY	
Risk-Aware Autonomous Decision Analysis under Cross-Domain Distribution Shift Using Uncertainty and Explanation Entropy	81
EMINE CEREN KARADEMİR, GÖKHAN UÇKAN, ARIFE ÇINAR AND EMRE ÇOMAK	
Approximation by Generalized Baskakov Operators Associated with Exponential Functions	82
GÜLSÜM ULUSOY ADA, MUSSA YOUSEF MOHAMED	
An Inertial Projection-Based Neurodynamic Approach for Image Restoration and Inpainting	83
RATTANAPORN WANGKEEREE, RABIAN WANGKEEREE, KASAMSUK UNGCHITTRAKOOL AND YIRGA ABEBE BELAY	
Weinstein Pseudo-Differential Operators on Torus	84
SANTOSH KUMAR UPADHYAY	

The Difference of Squares Formula and Quadratic Equations	85
SYRGAK KYDYRALIEV, ANARKUL URDALETOVA, ELENA BUROVA AND SALTANAT JAPAROVA	
Multi-State Performance Analysis of Benes Network	86
S. VISHAL VIGNESH , GUNDALA VIJAYALAKSHMI	
Convergence of the Generalised Picard-Mann Iteration for G-Monotone Non-Expansive Mappings in Banach Spaces with a Graph	87
DEEPA VIJAYASENAN, REKHA PANICKER AND RAHUL SHUKLA	
A New Gradient-Based Neurodynamic Network for Circular Cone Constrained Variational Inequality Problems	88
RATTANAPORN WANGKEEREE, RABIAN WANGKEEREE, YIRGA ABEBE BELAY, KASAMSUK UNGCHITTRAKOOL AND NUTT TANANIMIT	
A Projection Recurrent Neural Network for Circular Cone Constrained Variational Inequalities	89
RATTANAPORN WANGKEEREE, RABIAN WANGKEEREE, ALIREZA NAZEMI, YIRGA ABEBE BELAY AND KASAMSUK UNGCHITTRAKOOL	
The Numerical Range of the Generalized Inverse and Its Applications to Iterative Methods	90
GERALD WANJALA	
A New Approach to Decision Making Using Soft Covering Based Rough Sets by Concept of Soft Maximal Description	91
ZEHRA GÜZEL ERGÜL, KUBILAY YAŞAR	
Hyperbolic Pythagorean Hodograph Curves	93
AZIZ YAZLA, FATMA NUR DANIS	
Characterization of Some Classes of Compact Operators on Cesàro Difference Series Spaces	94
BUSE YUMUKTAY, CANAN HAZAR GÜLEÇ	
Modules with ss -Supplements in Their Cofinite Extensions	95
BAŞAK ZAMAN, EMINE ÖNAL KIR	

4th International Conference: Constructive Mathematical Analysis

Exponential Sampling Theory: Error Estimates via Mellin Distance

Carlo Bardaro

Department of Mathematics and Informatics, University of Perugia, Italy

`carlo.bardaro@unipg.it`

key-words: Mellin transform, Mellin distance, exponential sampling formula.

Abstract:

In the framework of Mellin–Lebesgue spaces, we introduce (see [1] and [2, Chapter 10]) a new notion of distance, using the inversion formula of the Mellin transform. This notion represents an efficient tool for estimating errors for some approximate basic formulae of Mellin analysis. In particular, we study the approximate versions of the exponential sampling formula, the reproducing kernel formula in Mellin setting, the Bernstein inequality for not necessarily band-limited functions in Mellin sense. It is proved that these estimates are asymptotically best possible and can be used also for obtaining characterizations of functional spaces in terms of the speed of convergence to zero of the errors. We study error estimates when the function f belongs to Mellin–Paley–Wiener spaces, Mellin inversion classes, Mellin–Lipschitz classes, Mellin–Sobolev spaces and Mellin–Hardy spaces, so obtaining a sort of hierarchy of these spaces in terms of the speed of convergence to 0 of the error.

References

- [1] Bardaro, C., Butzer, P.L., Mantellini, I. and Schmeisser, G. (2016). Mellin analysis and its basic associated metric—applications to sampling theory. *Analysis Math.*, 42(4), 297–321.
- [2] Bardaro, C., Butzer, P.L., Mantellini, I. and Schmeisser, G. (2025). *Mellin Analysis. Transform Theory and Applications*, Birkhauser, xxi + 487.

4th International Conference: Constructive Mathematical Analysis

Classes of Operators Related to Subnormal Operators

Raul E. Curto

Department of Mathematics, University of Iowa, Iowa City, Iowa, USA

raul-curto@uiowa.edu

key-words:

Abstract:

In this talk we lay the foundations for a new theory encompassing two natural extensions of the class of subnormal operators, namely the n -subnormal operators and the sub- n -normal operators, where $n \in \mathbb{N}$. We discuss inclusion relations among the above-mentioned classes and other related classes, e.g., n -quasinormal and quasi- n -normal operators. We show that sub- n -normality is stronger than n -subnormality, and produce a concrete example of a 3-subnormal operator which is not sub-2-normal. In [1], R.E. Curto, S.H. Lee and J. Yoon proved that if an operator T is subnormal, left-invertible, and such that T^2 is quasinormal, then T is quasinormal. In subsequent work, [3], P. Pietrzycki and J. Stochel improved this result by removing the assumption of left invertibility. In this talk we will consider suitable analogs of this result for the case of operators in the above-mentioned classes. In particular, we prove that the weight sequence of an n -quasinormal unilateral weighted shift must be periodic with period at most n .

The talk is based on joint work with Thankarajan Prasad, Nalanda University, India, which has recently appeared in print [2].

Research partially funded by U.S. National Science Foundation DMS-2247167.

References

- [1] Curto, R.E., Lee, S.H. and Yoon, J. (2020). Quasinormality of powers of commuting pairs of bounded operators. *J. Funct. Anal.*, 278, 108342.
- [2] Curto, R.E., Prasad, T. (2026). Classes of operators related to subnormal operators. *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.*, 120(2), 41.
- [3] Pietrzycki, P., Stochel, J. (2021). Subnormal n -th roots of quasinormal operators are quasinormal. *J. Funct. Anal.*, 280(12), 109001.

4th International Conference: Constructive Mathematical Analysis

Sampling Discretization of Integral Norms in Finite-Dimensional Spaces

Feng Dai

University of Alberta, Edmonton, Alberta, Canada

fdai@ualberta.ca

key-words: Marcinkiewicz discretization, random sampling, empirical L_p moments.

Abstract:

In this talk, I will present recent advancements in the Marcinkiewicz discretization problem using random sampling in finite-dimensional Spaces. The goal is to establish two-sided estimates for the integral norm of functions in the space via a finite sum of function values evaluated at randomly selected points that are independent of the individual functions in the space. The main challenge is to determine the “nearly” optimal number of random points required for the Marcinkiewicz discretization inequalities to hold with high probability.

References

- [1] Dai, F., Kosov, E. and Temlyakov, V. (2023). Some improved bounds in sampling discretization of integral norms. *J. Funct. Anal.*, 285, no. 4, Paper No. 109951.
- [2] Dai, F., Prymak, A., Shadrin, A., Temlyakov, V. and Tikhonov, S. (2021). Sampling discretization of integral norms. *Constr. Approx.*, 54, no. 3, 455–471.
- [3] Dai, F., Prymak, A., Shadrin, A., Temlyakov, V. and Tikhonov, S. (2021). Entropy numbers and Marcinkiewicz-type discretization. *J. Funct. Anal.*, 281, no. 6, Paper No. 109090.
- [4] Dai, F., Temlyakov, V. (2024). Random points are good for universal discretization. *J. Math. Anal. Appl.*, 529, no. 1, Paper No. 127570.
- [5] Dai, F., Kosov, E. and Temlyakov, V. (2026). A survey of sampling discretization of integral and uniform norms. arXiv preprint <https://arxiv.org/pdf/2603.02459> arXiv:2603.02459.

4th International Conference: Constructive Mathematical Analysis

Investigation of Physics-Informed Neural Networks for Solving the Allen-Cahn Equation

Adil Nauryzbay¹, Zhanars Abdiramanov^{1,2}, Esimkhan Zhumazhanov¹, Dastan Urnek¹
and Kazez Ertai¹

¹ Department of Mathematics and Mathematical Modeling, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan

² Institute of Information and Computational Technologies, Almaty, Kazakhstan

a.nauryzbai@abaiuniversity.edu.kz, zh.abdiramanov@abaiuniversity.edu.kz
yessimkhan.zhumazhanov@gmail.com, urnekdastan24@gmail.com

key-words: Physics-informed neural networks, Allen-Cahn equation, scalar auxiliary variable.

Abstract:

Physics-Informed Neural Networks (PINNs) offer a powerful deep learning framework for solving nonlinear partial differential equations. However, standard PINNs face significant challenges when applied to the Allen-Cahn equation, particularly concerning sharp interface resolution, spectral bias, and optimization stability during long-time integration. In this paper, we propose a novel Thermodynamically-Consistent Spectral-PINN architecture. To overcome optimization failures and Gibbs phenomena associated with classical spectral methods near sharp interfaces, we utilize Fourier Feature Positional Encoding. Furthermore, to guarantee strict energy dissipation without exacerbating the non-convexity of the loss landscape, we introduce a Scalar Auxiliary Variable (SAV) formulation into the PINN framework. We provide rigorous theoretical justification for the proposed architecture, including a proof that the modified energy functional is monotonically decreasing along solutions of the SAV-reformulated system. Preliminary numerical experiments are currently underway to validate the theoretical predictions.

References

- [1] Raissi, M., Perdikaris, P. and Karniadakis, G.E. (2019). Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear PDEs. *J. Comput. Phys.*, 378, 686-707.
- [2] Shen, J., Tang, T. and Wang, L.L. (2011). *Spectral Methods: Algorithms, Analysis and Applications*. Springer Science and Business Media, Berlin.
- [3] Shen, J., Xu, J. and Yang, J. (2018). The scalar auxiliary variable (SAV) approach for gradient flows. *J. Comput. Phys.*, 353, 407-416.
- [4] Shiyapov, K., Abdiramanov, Z., Issa, Z. and Zhumaseyitova, A. (2026). High-Order Spectral Scheme with Structure Maintenance and Fast Memory Algorithm for Nonlocal Nonlinear Diffusion Equations. *AppliedMath*, 6(4), 54.

4th International Conference: Constructive Mathematical Analysis

Modified Process Capability Index under Unit-Gumbel Distribution: Comparative Analysis of Point Estimators

Yunus Akdogan¹, Tenzile Erbayram¹

¹ Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

yakdogan@selcuk.edu.tr, tenzile.erbayram@selcuk.edu.tr

key-words: Process capability index, unit distribution, Monte Carlo simulation.

Abstract:

Process capability indices are fundamental statistical indicators that quantitatively measure a process's capacity to produce output within defined specification limits and are of strategic importance in quality control processes. This study presents a specialized C_{GPK} index based on the unit Gumbel distribution to determine the process capability index for unit distributions, a limited area in the literature. In this study, point estimates are obtained for the C_{GPK} index using different statistical methods. To determine the effectiveness of the obtained estimators, the performance measures used bias, mean squared errors, and mean relative errors are compared through a comprehensive simulation study. The simulation results reveal the effectiveness of the estimation methods under different sample sizes and parameter combinations. Theoretical findings, applications conducted on various datasets defined within the unit range (0,1), and the contribution of this index to the literature and its practical advantages in process performance evaluation have been proven.

References

- [1] Arslan, T. (2023). A new family of unit-distributions: Definition, properties and applications. TWMS Journal of Applied and Engineering Mathematics.
- [2] Kashif, M., Aslam, M., Rao, G. S., AL-Marshadi, A. H. and Jun, C. H. (2017). Bootstrap confidence intervals of the modified process capability index for Weibull distribution. Arabian Journal for Science and Engineering, 42(11), 4565-4573.

4th International Conference: Constructive Mathematical Analysis

Geometric Analysis of Magnetic Trajectories of Charged Particles and Their Relations with Associated Curves

Hüda Nur Akgül¹, Muhammed T. Sariaydin¹

¹ Department of Mathematics, Faculty of Science, Selcuk University, Konya, Türkiye

hnakgul17@gmail.com, talatsariaydin@gmail.com

key-words: Magnetic Trajectory, associated curves, Lorentz equation.

Abstract:

This paper focuses on the geometric analysis of trajectories of charged particles moving under the influence of magnetic fields. The primary aim is to reveal the relationships between these trajectories and special curves in differential geometry. In this context, magnetic curves are examined in relation to well-known curve pairs, and several of their fundamental geometric properties are established. Finally, illustrative examples are presented to support the theoretical results.

References

- [1] As, E., Sarıoğlu, A. (2014). On the Bishop curvatures of involute-evolute curve couple in. Int. J. Phys. Sci, 9(7), 140-145.
- [2] Bilici, M., Çalışkan, M. (2009). On the Involute of the Spacelike Curve with a Timelike Binormal in Minkowski 3-Space. In International Mathematical Forum, 4(31), 1497-1509.
- [3] Bozkurt, Z., Gök, I., Yaylı, Y. and Ekmekci, F. N. (2014). A new approach for magnetic curves in 3D Riemannian manifolds. Journal of Mathematical Physics, 55(5).
- [4] Izumiya, S., Takeuchi, N. (2002). Generic properties of helices and Bertrand curves. Journal of Geometry, 74(1), 97-109.
- [5] Orbay, K., Kasap, E. (2009). On Mannheim partner curves in E^3 International Journal of Physical Sciences, 4(5), 261-264.
- [6] Sariaydin, M. T., Yazla, A. (2023). Associated curves of charged particle moving with the effect of magnetic field. Commun. Korean Math. Soc. 38(2): 589-598.

4th International Conference: Constructive Mathematical Analysis

OMNIAI: AI-Powered Chatbot-Based Appointment Management System

Muhammet İkbāl Akgül¹, Gökhan Uçkan¹ and Emre Çomak¹

¹ Department of Mathematics, Pamukkale University, Denizli, Türkiye

muhammedakg23@gmail.com, guckan@pau.edu.tr, ecomak@pau.edu.tr.

key-words: Artificial intelligence, cloud computing, microservices.

Abstract:

The rapid digitalization of customer-business interactions has highlighted the inefficiencies of traditional, manual appointment management systems, which are often prone to high latency, human error, and limited operational hours. To resolve these challenges, this study presents OmniAI, an artificial intelligence-powered, chatbot-based appointment management system designed to fully automate the reservation process and optimize administrative workflows. The proposed method leverages a microservices architecture, predominantly developed using the Go programming language, to ensure high scalability and modularity. The system interfaces directly with end-users via the WhatsApp messaging platform using the Twilio API. To handle concurrent requests and maintain high performance under heavy traffic, asynchronous message queuing is implemented via RabbitMQ. The core intelligence of OmniAI is driven by an AI Service integrated with OpenAI's GPT-4o model. This is significantly enhanced by a Retrieval-Augmented Generation (RAG) module, which dynamically queries a PostgreSQL-backed corporate knowledge base. This integration enables the chatbot to fetch context-specific data—such as working hours, branch details, and service descriptions—facilitating highly accurate intent analysis and context-aware natural language responses. Furthermore, the system includes a comprehensive frontend dashboard built with React and TypeScript, allowing business administrators to monitor daily reservations, manage user roles with JWT-based authentication, and track AI performance analytics in real-time. The entire ecosystem is containerized using Docker and securely routed through a Traefik API gateway, ensuring isolated and portable deployments. The implementation results demonstrate that OmniAI substantially decreases average response times and reservation errors compared to traditional methods, while maintaining a high rate of successful automated bookings. In conclusion, the integration of advanced Large Language Models with RAG in a microservices framework provides a robust, 24/7 automated booking experience, significantly reducing administrative overhead and improving overall customer satisfaction.

References

- [1] Dragoni, N. et al. (2017). Microservices: How To Make Your Application Scale. Present and Ulterior Software Engineering. Cham: Springer, 95-104.
- [2] Lewis, P. et al. (2020). Retrieval-augmented generation for knowledge-intensive NLP tasks. Advances in Neural Information Processing Systems, 33, 9459-9474.
- [3] Merkel, D. (2014). Docker: lightweight Linux containers for consistent development and deployment. Linux Journal, (239), 2.
- [4] Newman, S. (2021). Building Microservices: Designing Fine-Grained Systems, 2nd ed. Sebastopol, CA: O'Reilly Media.
- [5] OpenAI, (2023). GPT-4 Technical Report. arXiv preprint. arXiv:2303.08774.

4th International Conference: Constructive Mathematical Analysis

On Integral Curves Generated by PAF in Euclidean 3-Space

Selen Akpınar¹, Muhammed T. Sariaydin¹

¹ Department of Mathematics, Faculty of Science, Selcuk University, Konya, Türkiye

selenakpinar42@gmail.com, talatsariaydin@gmail.com

key-words: Integral curves, associated curves, Euclidean space.

Abstract:

This study, we introduces a family of integral curves associated with a curve according to PAF in Euclidean 3-space. The curvature relations among these curves are analyzed comprehensively. Moreover, the developed approach offers constructive procedures for obtaining certain special curves. To illustrate and validate the theoretical findings, several representative examples are included.

References

- [1] Duldul, M., Bulbul, Z. (2023). Integral curves connected with a framed curve in 3-space. Honam Mathematical J., 45(1), 130-145.
- [2] Kandilci, G. T., Ekici, C. and Dede, M. (2023). Directional Bertrand Curves in Minkowski Space. Innovative Research in Natural Science and Mathematics, 15.
- [3] Korpınar, T., Sariaydin, M. T. and Turhan, E. (2013). Associated curves according to Bishop frame in Euclidean 3-space. Advanced Modeling and Optimization, 15(3), 713-717.
- [4] Şahiner, B. (2019). Direction curves of tangent indicatrix of a curve. Applied Mathematics and Computation, 343, 273-284.
- [5] Doğan Yazıcı, B., Özkaldı Karakuş, S. and Tosun, M. (2022). Framed Curves and Their Applications Based on a New Differential Equation. International Electronic Journal of Geometry, 15(1), 47-56.

4th International Conference: Constructive Mathematical Analysis

Fixed Point Approximation Schemes for Wasserstein Gradient Flows in Generative Modeling

Osman Alagöz¹, Emrah Haspolat¹

¹ Department of Mathematics, Bilecik Şeyh Edebali University, Bilecik, Türkiye

osman.alagoz@bilecik.edu.tr, emrah.haspolat@bilecik.edu.tr

Keywords: Wasserstein gradient flow, generative modeling, proximal operator.

Abstract:

We study a conditional fixed point framework for a particle-based Wasserstein proximal sampler. Motivated by the Jordan–Kinderlehrer–Otto scheme, we regard a Wasserstein proximal step as a self-map and define Picard, Mann and Ishikawa updates through selected Wasserstein geodesics. Under explicit contractivity and selected-geodesic metric-convexity assumptions, we prove convergence and perturbation estimates. In addition, we identify the empirical quadratic proximal step induced by optimal matching, which provides a controlled common-coupling particle surrogate where the assumptions can be checked. Numerical experiments on two-dimensional targets compare fixed-target, mini-batch, and resampled-target regimes. Picard/JKO is the strongest method in deterministic fixed-target tests, while relaxation, especially Mann relaxation, can reduce the final empirical W_2 error when the target batch is resampled. The results are intended as a stability analysis for this particle surrogate, not as a global contractivity theorem for general Wasserstein proximal maps or as a large-scale generative modeling claim.

References

- [1] Cuturi, M. (2013). Sinkhorn distances: Lightspeed computation of optimal transport. *Advances in Neural Information Processing Systems*, 26, 2292–2300.
- [2] Karras, T., Aittala, M., Aila, T. and Laine, S. (2022). Elucidating the design space of diffusion-based generative models. *Advances in Neural Information Processing Systems*, 35, 26565–26577.
- [3] Lipman, Y., Chen, R.T.Q., Ben-Hamu, H., Nickel, M. and Le, M. (2023). Flow matching for generative modeling. *International Conference on Learning Representations*.

4th International Conference: Constructive Mathematical Analysis

Two Dimensional Fuzzy Differential Transform Method for Solving of Two Dimensional Nonlinear Fuzzy Volterra Integral Equations

Artan F. Alidema¹, Frederik Dara²

¹ Department of Mathematics, University of Prishtina, Prishtina, Kosovo

² Department of Applied Mathematics, University of Tirana, Tirane, Albania

artan.alidema@uni-pr.edu, frederik.dara@fshn.edu.al

key-words: Fuzzy differential transform method, two dimensional fuzzy Volterra integral equations, fuzzy calculus.

Abstract:

This study applies the two-dimensional fuzzy differential transform method (2D-FDTM) to solve two-dimensional fuzzy Volterra integral equations (2D-FVIEs). The fundamental properties of the method are established, and illustrative examples are provided to verify its efficiency and accuracy. The results confirm that the proposed approach offers a reliable analytical tool for handling fuzzy Volterra integral equations in two dimensions.

References

- [1] Georgieva A., Alidema A. (2019). Convergence of homotopy perturbation method for solving of two-dimensional fuzzy Volterra functional integral equations. AIP Conf Proc., 1910, 050009.
- [2] Khaji, R., Yaseen, F. and Khalil, R. I. (2024). Solution of fuzzy Volterra integral equations using composite method. Adv Math Models Appl., 9(1), 50–60.
- [3] Mikaeilvand N, Khakrangin S. (2012). Solving fuzzy partial differential equations by fuzzy two-dimensional differential transform method. Neural Computing and Applications, 135, 307–312.
- [4] Puri M. L., Ralescu D. A. (1983). Differential of fuzzy function. J Math Anal Appl., 91, 552–558.
- [5] Sadatrasoul S., Ezzati R. (2016). Numerical solution of two-dimensional nonlinear Hammerstein fuzzy integral equations based on optimal fuzzy quadrature formula. J Comput Appl Math, 292, 430–446.
- [6] Sadatrasoul S., Ezzati R. (2014). Quadrature rules and iterative method for numerical solution of two-dimensional fuzzy integral equations. Abstr Appl Anal., 18 pages.
- [7] Salhasour S., Allahviranloo T. (2013). Application of fuzzy differential transform method for solving fuzzy integral equations. Applied Mathematical Modelling, 37(3), 1016–1027.
- [8] Tari A, Shahmorad S. (2011). Differential transform method for the system of two-dimensional nonlinear Volterra integro-differential equations. Computers and Mathematics with Applications, 61(9), 2621–2629.
- [9] Xavier D., Behera D. and Xavier S. D. (2025). Double parametric based solution of fuzzy Volterra integral equations with separable type kernels. J. Adv. Appl. Comput. Math., 15(2), 88–98.

4th International Conference: Constructive Mathematical Analysis

On the Approximation of Borel Derivatives by Nonlinear Szász Mirakyan Durrmeyer Operators

Hüseyin Erhan Altın

Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

erhanaltin@ibu.edu.tr

key-words: Borel derivative, nonlinear operator, pointwise convergence.

Abstract:

In the present study, we consider a sequence of nonlinear Szász-Mirakyan-Durrmeyer operators (NS_n) of the form

$$(NS_n)(f; x) = \int_0^{\infty} K_n(x, t, f(t)) dt, \quad x \in [0, \infty), \quad n \in \mathbb{N},$$

acting on bounded functions on every finite subinterval of $[0, \infty)$ where $K_n(x, t, u)$ satisfy some suitable assumptions.

We give some approximation results with regard to the convergence of the operators NS_n to right, left, and symmetric Borel differentiable functions.

Our study extend some of the previous results about the linear integral operators and also aims to bring a new perspective to research in approximation with nonlinear operators by introducing a different concept of a derivative.

References

- [1] Altın, H. E., Karslı, H. (2021). Some approximation properties of a nonlinear Szász-Mirakyan-Durrmeyer operator. *Analele Universitatii Oradea Fasc. Matematica*, Tom XXVIII, (1), 105-114.
- [2] Bardaro, C., Musielak, J. and Vinti, G. (2003). Nonlinear integral operators and applications. *De Gruyter Series in Nonlinear Analysis and Applications*, 9, xii - 201.
- [3] Ibikli, E. (2001). Approximation of Borel derivatives of functions by singular integrals. *Trans. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci.* 21(4), *Math. Mech.*, 72-76.
- [4] Karslı, H., Tiryaki, I. U. (2021). Approximation of Borel derivatives of functions via non-linear singular integral operators. *Palestine Journal of Mathematics*. 10(2), 673-684.
- [5] Musielak, J. (1983). On some approximation problems in modular spaces. In *Constructive Function Theory 1981 (Proc. Int. Conf., Varna, June 01-05, 1981, pp. 455-461)*, Publ. House Bulgarian Acad. Sciences.
- [6] Sahai, A., Prasad, G. (1993). On the rate of convergence for modified Szász-Mirakyan operators on functions of bounded variation. *Publ. Inst. Math. (beograd) (N.S.)* 53(67), 73-80.

4th International Conference: Constructive Mathematical Analysis

Almost Automorphic and Pseudo-Almost Automorphic Solutions: A Unified Theory with Applications to Classical and Fractional Evolution Systems

Dhaou Lassoued^{1,2}, Jehad Alzabut^{3,4}

¹ Faculty of Sciences of Gabès, Department of Mathematics, Cité Erriadh Gabès, Tunisia

² SAMM EA4543 Laboratory, University Paris 1 Panthéon-Sorbonne, Paris cedex 13, France

³ Department of Mathematics and Sciences, Prince Sultan University, Riyadh, Saudi Arabia

⁴ Department of Industrial Engineering, OSTIM Technical University, Ankara, Türkiye

dhaou06@gmail.com, jalzabut@psu.edu.sa

key-words: Almost automorphic functions, pseudo almost automorphic functions, parabolic equations.

Abstract:

This paper studies almost automorphic and pseudo-almost automorphic mild solutions for a class of non-linear evolution equations in Banach spaces, covering both classical and fractional frameworks. A unified approach is developed based on semigroup and resolvent theory, together with fixed-point techniques, to establish existence and uniqueness results for semilinear equations generated by exponentially stable C_0 -semigroups and their fractional counterparts. The novelty lies in providing a coherent framework that simultaneously treats classical and fractional dynamics while incorporating both almost automorphic and pseudo-almost automorphic structures. The results extend several known contributions in the literature. Applications are presented to parabolic integrodifferential equations with memory, the Schrödinger–Poisson system, and classical and fractional Navier–Stokes equations, including a concrete example illustrating the pseudo-almost automorphic behavior of the forcing term.

References

- [1] Bochner, S. (1962). A new approach to almost periodicity. Proc. Natl. Acad. Sci. USA, 48(12), 2039–2043.
- [2] Bochner, S. (1964). Continuous mappings of almost automorphic and almost periodic functions. Proc. Natl. Acad. Sci. USA, 52, 907–910.
- [3] Levitan, B.N., Zhikov, V.V. (1982). Almost-Periodic Functions and Differential Equations. Cambridge University Press.
- [4] Veech, W. A. (1963). Almost automorphic functions. Proc. Natl. Acad. Sci. USA, 49, 462–464.
- [5] Veech, W. A. (1965). Almost automorphic functions on groups. Amer. J. Math., 87, 719–751.

4th International Conference: Constructive Mathematical Analysis

Numerical Study on a Class of Higher-Order Singular Polytropic Equations

Djihene Arafa¹, Nouria Arar²

¹ Laboratory of Applied Mathematics and Didactics, Department of Mathematics, Higher Normal School Assia Djebar, Constantine, Algeria

² Department of Mathematics, University Constantine 1 - Brothers Mentouri, Constantine, Algeria

arafa.djihene@ensc.dz, arar.nouria@umc.edu.dz

key-words: Collocation method, fractional Legendre polynomials, operational matrices.

Abstract:

In this work, we address the numerical approximation of a class of higher-order singular polytropic equations (SPEs) with variable coefficients, written in the general form [2]

$$x^{-\eta} D^{\beta}(x^{\eta} D^{\gamma} v(x)) = \omega(x, v(x)), \quad (1)$$

where $v(x)$ is the unknown function, $\eta > 0$ is a real shape parameter, $\omega(x, v(x))$ is a given (possibly nonlinear) source term, and β, γ are Caputo fractional differentiation orders, with $\beta + \gamma$ defining the total order of (1).

We develop a collocation scheme based on shifted fractional-order Legendre polynomials $\mathcal{L}_n^{(\alpha)}(x)$ on $[0, 1]$. Operational matrices for fractional integration and product are constructed in this basis, reducing (1) to a linear or nonlinear algebraic system for the spectral coefficients. A rigorous error bound confirms the reliability and convergence of the proposed approach. Numerical experiments on fourth- and fifth-order test problems show that an appropriate choice of α reduces the absolute error by several orders of magnitude while requiring fewer basis functions than existing techniques such as the Haar wavelet method [1].

References

- [1] Albalawi, K.S., Kumar, A., Alkahtani, B.S. and Goswami, P. (2023). Solving general fractional differential equations using Haar wavelet collocation method. *Fractal and Fractional*, 7(8).
- [2] Ali, K. K., Mehanna, M. S., Abdelrahman, M. I. and Shaalan, M. A. (2022). Analytical and Numerical solutions for fourth order Lane–Emden–Fowler equation. *Partial Differential Equations in Applied Mathematics*, 6, 100430.
- [3] Podlubny, I. (1999). *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*. Elsevier Science.
- [4] Qu, H., Yang, X., and She, Z. (2020). Left and right-shifted fractional Legendre functions with an application for fractional differential equations. *Adv. Math. Phys.*, 2020(1), 6036417.

4th International Conference: Constructive Mathematical Analysis

On a New Modification of the Mellin-Gauss-Weierstrass Operators in the Weighted Mellin-Lebesgue Spaces

Ali Aral

Department of Mathematics, Kırıkkale University, Kırıkkale, Türkiye

aliaral73@yahoo.com

key-words: Logarithmic weighted modulus of continuity, logarithmic moments, weighted Mellin-Lebesgue spaces.

Abstract:

In this presentation, we establish a quantitative convergence result for a new generalization of Mellin-Gauss-Weierstrass operators preserving logarithmic functions, using the logarithmic weighted modulus of continuity. We derive the logarithmic moments of the modified operators and express a Voronovskaya-type theorem. Furthermore, we obtain the rate of convergence and show global smoothness preservation property in weighted Mellin-Lebesgue spaces of Lebesgue measurable functions.

References

- [1] Aral, A. (2023). On a new approach in the space of measurable functions. *Constr. Math. Anal.*, 6(4), 237-248.
- [2] Aral, A., Acar, T., Kursun, S. (2022). Generalized Kantorovich forms of exponential sampling series. *Anal. Math. Phys.*, 12(2), No. 50.
- [3] Aral, A., Bardaro, C., Mantellini, I., Ozsarac, F. (2025). A note on the Mellin-Fejer kernels. *Results Math.*, 80(3), No:80.
- [4] Bardaro, C., Mantellini, I. (2011). Asymptotic behaviour of Mellin-Fejer convolution operators. *East J. Approx.*, 17(2), 181-201.
- [5] Bardaro, C., Mantellini, I. (2014). On Mellin convolution operators: a direct approach to the asymptotic formulae. *Integral Transf. Spec. Funct.*, 25(3), 182-195.
- [6] Butzer, P. L., Jansche, S. (1997). A direct approach to the Mellin transform. *J. Fourier Anal. Appl.*, 3(4), 325-376.
- [7] Mamedov, R. G. (1991). *The Mellin Transform and Approximation Theory*, Elm, Baku.
- [8] Ozsarac, F. (2024). On the Mellin-Gauss-Weierstrass operators in the Mellin-Lebesgue spaces,. *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 73(3), 630-640.
- [9] Topuz, C., Ozsarac, F., Aral, A. (2024). On the generalized Mellin integral operators. *Demonstr. Math.*, 57(1), pp. 20230133.

4th International Conference: Constructive Mathematical Analysis

On Some Matrix Transformations of Sequences of Function

Ali Arpacioğlu^{1,2}, Tuncay Tunç³

¹ Human Resources Management Program, Tarsus University, Mersin, Türkiye

² Institute of Science, Department of Mathematics, Mersin University, Mersin, Türkiye

³ Department of Mathematics, Mersin University, Mersin, Türkiye

aliarpacioglu@tarsus.edu.tr, ttunc@mersin.edu.tr

key-words: Cesàro matrix, function sequences, Zweier matrix.

Abstract:

In mathematical analysis, the convergence of function sequences and series is a fundamental concept in both theoretical and applied mathematics. Different types of convergence of function sequences and their behavior in function spaces are significant area of research analysis and functional analysis. In recent years, research has focused on alpha convergence, one of the important types of convergence of function sequences, as defined below, and its generalizations. Let (X, d) be a metric space. Let $(f_n) \in \mathfrak{F}(\mathbb{R}^X)$, $f \in \mathbb{R}^X$, and $x_0 \in X$ be given. The sequence of functions (f_n) is said to be α -convergent to the function f at the point x_0 if $f_n(x_n) \rightarrow f(x_0)$ for every sequence $(x_n) \subset X$ such that $x_n \rightarrow x_0$ [1].

In this study, we investigate the convergence properties of real-valued function sequences defined on a metric space that are pointwise convergent, uniformly convergent, alpha-convergent, and uniformly alpha-convergent under Zweier and Cesàro matrix transformations. We will also examine the behavior of sequences with pointwise and uniform exhaustiveness properties under these summability matrices.

References

- [1] Courant, R. (1914). Ueber eine Eigenschaft der Abbildungsfunktionen bei konformer Abbildung. Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, 101-109.

4th International Conference: Constructive Mathematical Analysis

A Neurodynamical Neural Network Approach for Solving Support Vector Machine Problems with Generalized Pinball Loss Functions via Hyperbolic Tangent Smoothing Technique

Rattanaorn Wangkeeree¹, Rabian Wangkeeree¹, Yirga Abebe Belay^{1,2} and Kasamsuk Ungchittrakool¹

¹ Department of Mathematics, Naresuan University, Phitsanulok, Thailand

² Department of Mathematics, Aksum University, Axum, Ethiopia

rattanaornw@nu.ac.th, rabianw@nu.ac.th, yirga2006@gmail.com, kasamsuku@nu.ac.th

key-words: Neural networks, support vector machines.

Abstract:

In this paper, we propose a neurodynamical neural network approach for solving support vector machine problems with generalized pinball loss function. First, we use the hyperbolic tangent smoothing technique to find a smooth approximation of the generalized pinball loss associated with the SVM problem. Then we propose a neurodynamical neural network model to solve the given support vector machine problem with smooth objective function. We prove the equivalence between the set of optimal solutions the smoothed optimization problem and the set of equilibrium points of the proposed model. Then we analyze the existence and uniqueness of solutions, stability, and convergence of the method. Next we conduct numerical simulations both on synthetic and UCI datasets to show the effectiveness of the method. We evaluate the performance of the model using the accuracy, Matthews Correlation Coefficient (MCC), and F1-score metrics. Finally, we compare our model against existing models of solving support vector machine problems with pinball loss functions. We conduct the Friedman test statistical analysis followed by the Nemenyi post-hoc analysis to examine the statistical difference of the proposed model against the existing methods.

References

- [1] Huang, X., Shi, L. and Suykens, J. A. K. (2014). Support vector machine classifier with pinball loss. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 35(6), 984–997.
- [2] Makmuang, D., Ratiphaphongthon, W. and Wangkeeree, R. (2023). Smooth support vector machine with generalized pinball loss for pattern classification. *The Journal of Supercomputing*, 79(11), 11684–11706.
- [3] Panup, W., Ratipapongton, W. and Wangkeeree, R. (2022). A novel twin support vector machine with generalized pinball loss function for pattern classification. *Symmetry*, 14(2), Article 289.
- [4] Reshma, R., Khemchandani, R., Pal, A. and Chandra, S. (2018). Generalized pinball loss SVMs. *Neurocomputing*, 322, 10–22.
- [5] Suppalap, S., Makmuang, D., Damminsud, V. and Wangkeeree, R. (2025). Smooth support vector machine with rescaled generalized pinball loss for classification. *The Journal of Supercomputing*, 81(1), Article 217.
- [6] Wangkeeree, R., Wangkeeree, R., Belay, Y. A., Ungchittrakool, K., and Nazemi, A. (2026). Neurodynamical neural network method for solving support vector machines with epigraph reformulated generalized pinball loss function. *Journal of Computational and Applied Mathematics*, Article 117754.

4th International Conference: Constructive Mathematical Analysis

Where/Why Do We Use the Bivariate/Multivariate Positive Linear/Nonlinear Operators?

Murat Bodur

Department of Engineering Basic Sciences, Konya Technical University, Konya, Türkiye

mbodur@ktun.edu.tr

key-words: Positive linear/nonlinear operators, computer-aided geometric design, neural networks.

Abstract:

Approximation theory is a fundamental branch of mathematical analysis concerned with the representation of complex functions by simpler, more computationally tractable ones, such as polynomials, splines, or rational functions. In this presentation, we investigate the theoretical motivations and practical implementations of bivariate and multivariate positive linear and nonlinear operators within the framework of approximation theory.

Acknowledgements: The author has been supported by the Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country.

References

- [1] Altomare, F., Campiti, M. (1994). Korovkin-Type Approximation Theory and its Applications. Walter de Gruyter, Berlin.
- [2] DeVore, R. A. (1998). Nonlinear Approximation. Acta Numerica, Cambridge University Press, 7, 51–150.
- [3] Farin, G. (2014). Curves and Surfaces for Computer-Aided Geometric Design: A Practical Guide. Academic Press.
- [4] Pinkus, A. (1999). Approximation Theory of the MLP Model in Neural Networks. Acta Numerica, 8, 143–195.

4th International Conference: Constructive Mathematical Analysis

Existence of Solutions for a Fractional $p(x, \cdot)$ -Laplacian Problem with Neumann Boundary Conditions on an Exterior Domain

Abdelkrim Barbara¹, Ahmed Bousmaha¹ and Mohammed Shimi²

¹ Laboratory of Modeling, Applied Mathematics, and Intelligent Systems, Sidi Mohamed Ben Abdellah University, Fez, Morocco

² Laboratory of Mathematics and Applications to Engineering Sciences, Sidi Mohamed Ben Abdellah University, Fez, Morocco

abdelkrim.barbara@usmba.ac.ma, ahmed.bousmaha@usmba.ac.ma, mohammed.shimi2@usmba.ac.ma

key-words: Fractional $p(x, \cdot)$ -laplacian, Nehari manifold, nonlocal nonlinear boundary condition, exterior domain.

Abstract:

This paper addresses the existence of non-negative solutions for a fractional $p(x, \cdot)$ -Laplacian problem with nonlocal nonlinear boundary conditions on an exterior domain. The study is conducted using the Nehari manifold approach, which is the primary instrument for establishing the key results.

References

- [1] Chen, C., Zhang, Q. (2012). Multiple solutions of a quasilinear elliptic problem involving nonlinear boundary condition on exterior domain. Math Methods Appl Sci, 35, 782–794.
- [2] Zhao, L., Li, A. and Su, J. (2012). Existence and multiplicity results for quasilinear elliptic exterior problems with nonlinear boundary conditions. Nonlinear Anal, 75, 2520–2533.

4th International Conference: Constructive Mathematical Analysis

Magic Squares: From Ancient Aesthetics to a New Recursive Construction Method

Syrgak Kydyraliev¹, Anarkul Urdaletova², Elena Burova¹ and Saltanat Japárova³

¹ Department of Applied Mathematics and Informatics, American University of Central Asia, Bishkek, Kyrgyz Republic

² Department of Mathematics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyz Republic

³ Department of Mathematics, Issyk-Kul State University named after Kasym Tynystanov, Karakol, Kyrgyz Republic

kydyraliev_s@auca.kg, anarkul.urdaletova@manas.edu.kg, burova_e@auca.kg, japárova@iksu.kg

key-words: Magic squares, recursive construction, matrix theory.

Abstract:

The history of solving quadratic equations stretches back over four thousand years to Ancient Babylon, where scribes calculated land areas and astronomical cycles using rules remarkably similar to our own. However, most cuneiform texts provide only "recipes"—step-by-step instructions without explaining the underlying logic. Paradoxically, modern mathematics education often mirrors this ancient approach, favoring universal formulas (the "standard recipe") while neglecting more elegant and efficient alternatives. This study demonstrates that a wide range of problems typically solved via standard quadratic equations can be addressed more effectively through the direct application of the difference of squares formula.

We propose a specific algorithm based on this approach, highlighting its relevance for mathematical modeling. Many real-world problems, when initially translated into mathematical language, naturally take a form that is ideally suited for the difference of squares method before they are forcibly reduced to a standard quadratic form. By revisiting techniques once championed by Diophantus, Al-Khwarizmi, and Fibonacci—methods largely overshadowed since the 17th century—this work offers a streamlined alternative to the "general recipe," enhancing both computational efficiency and conceptual clarity in the classroom.

References

- [1] Kydyraliev, S. K., Urdaletova, A. B. (2020). Fashionable formula. Vestnik KRSU, 20(4), 3-9.
- [2] Kydyraliev, S. K., Urdaletova, A., Japárova, S. N., and Burova, E. (2025). From Diophantus and Al-Khwarizmi to the Modern Times. Tech Fusion in Business and Society. Studies in Systems, Decision and Control, 233, 335-339.
- [3] Kydyraliev, S. K., Urdaletova, A. B. (2017). Living Mathematics. 7th grade. Bishkek: Solution.

4th International Conference: Constructive Mathematical Analysis

Eigenvalue Expansions for a Conformable Sturm–Liouville Operator with Mixed Boundary Conditions

Olgun Cabri

Department of Business Management, Artvin Çoruh University, Artvin, Türkiye

olguncabri@artvin.edu.tr

key-words: Conformable derivative, Sturm–Liouville operator, perturbation theory.

Abstract:

In this paper, we study the conformable Sturm–Liouville problem

$$-D^{2\alpha}y(x) = \lambda y(x), \quad y(0) = 0, \quad D^\alpha y(1) = 0, \quad (1)$$

on the unit interval, where $D^\alpha y(x) = x^{1-\alpha}y'(x)$ denotes the conformable derivative [2]. A Dirac delta perturbation of the form $V(x) = \delta(x - a)$ is introduced, and the corresponding first- and second-order eigenvalue corrections are derived. We analyze the critical points of the first-order correction with respect to the perturbation point a and the conformable order α . As the two endpoints are not symmetric, we study the behaviour of the spectrum under perturbations near the Neumann end $x = 1$. Moreover, the cumulative response over a right subinterval is obtained in closed form.

References

- [1] Anderson, D. R., Camrud, E. and Ulness, D. J. (2019). On the nature of the conformable derivative and its applications to physics. *J. Fract. Calc. Appl.*, 10(2), 92–135.
- [2] Khalil, R., Al Horani, M., Yousef A. and Sababheh, M. (2014). A new definition of fractional derivative. *J. Comput. Appl. Math.*, 264, 65–70.

4th International Conference: Constructive Mathematical Analysis

Appell-Vietoris Polynomials and Their Zeros

Isabel Cação¹, Maria Irene Falcão², Helmuth Robert Malonek¹ and Graça Tomaz³

¹ Department of Mathematics, University of Aveiro, Aveiro, Portugal

² Centro de Matemática, Universidade do Minho, Braga, Portugal

³ Polytechnic of Guarda, Guarda, Portugal

isabel.cacao@ua.pt, mif@math.uminho.pt, hrmalon@ua.pt, gtomaz@ipg.pt

key-words: Vietoris number sequence, Appell-Vietoris polynomials, zero bounds.

Abstract:

The Vietoris numbers form a sequence of pairwise repeated rational numbers that originally arose in the study of the positivity of certain trigonometric sums (cf. [2]). In this talk, we consider the class of Appell polynomials associated with the Vietoris sequence, introduced in [1], and investigate several of their structural properties. In particular, we derive lower and upper bounds for their zeros in the complex plane.

References

- [1] Cação, I., Falcão, M.I., Malonek, H.R., Miranda, F. and Tomaz, G. (2024). On Appell-Vietoris polynomials. In: Gervasi, O., et al. (eds.) ICCSA 2024. LNCS, (14815), 302–316. Springer, Cham.
- [2] Vietoris, L. (1958). Über das Vorzeichen gewisser trigonometrischer Summen, Sitzungsber. Österr. Akad. Wiss 167, 125–135.

4th International Conference: Constructive Mathematical Analysis

Approximation Processes on Minimal Surfaces

Michele Campiti

Department of Mathematics and Physics “E. De Giorgi”, University of Salento, Lecce, Italy

michele.campiti@unisalento.it

key-words: Minimal surfaces, non-linear Bernstein-Schnabl operators, Voronovskaja formula.

Abstract:

We point out the non-additive potential structures and continuous-time evolutionary processes generated by a class of non-linear Bernstein-Schnabl type operators on minimal surfaces. Through a generalized Voronovskaja-type formula, we compute the uniform limit and the connection with a degenerate second-order elliptic differential operator whose coefficients are explicitly governed by the intrinsic geometry. For a fixed initial datum f , the evolution trajectory remains strictly constrained to the Riemannian metric g_f spanned by that datum. This allows us to treat the non-linear iteration identically to a sequence of positive linear contractions. This extends to a much wider context the results in [1, 2].

References

- [1] Altomare, F. (1989). Limit semigroups of Bernstein-Schnabl operators associated with positive projections. *Annali Sc. Norm. Sup. Pisa*, 16(2), 259–279.
- [2] Altomare, F., Campiti, M. (1994). *Korovkin-type Approximation Theory and its Applications*. De Gruyter Studies in Mathematics 17 W. De Gruyter, Berlin-New York.

4th International Conference: Constructive Mathematical Analysis

Statistical Inference of a New Four-Parameter Marshall Olkin Distribution: Comprehensive Simulation and Measurement Applications

Erdem Cankut¹, Kadir Karakaya¹

¹ Department of Statistics, Faculty of Science, Selçuk University, Konya, Türkiye

erdem.cankut@selcuk.edu.tr, kkarakaya@selcuk.edu.tr

key-words: Marshall Olkin family, Monte Carlo simulation, measurement datasets.

Abstract:

In this study, a new flexible four-parameter Marshall–Olkin Fréchet Power Rayleigh (MOFPR) distribution is introduced, which is constructed by applying the Marshall–Olkin [2] family of distributions to the Fréchet Power Rayleigh [1] model, and its statistical properties are examined. The fundamental mathematical and statistical characteristics of the proposed model, including the probability density function, cumulative distribution function, hazard rate function, and quantile function, are derived. To illustrate the model's versatility, the behavior of the probability density function and cumulative distribution function are analyzed through graphical representations under various parameter configurations. A significant contribution of this research is the comprehensive estimation framework, where twelve different estimation methods—including maximum likelihood, least squares, weighted least squares, Anderson–Darling, and Cramér–von Mises, among others—are employed to estimate the unknown parameters. The finite-sample performance of these estimators is assessed through an extensive Monte Carlo simulation study involving 5,000 replications. The efficiency of the estimators is compared based on several performance metrics, such as bias, mean squared error, mean relative error, and average absolute bias. Furthermore, the practical utility of the proposed distribution is demonstrated using two real measurement datasets. The modeling capability of the new distribution is evaluated via goodness-of-fit statistics, including Kolmogorov–Smirnov, Anderson–Darling, and Cramér–von Mises tests. Comparative results indicate that the proposed model provides a superior fit over several distributions in the literature, such as the Weibull, gamma, exponential, Fréchet, and Rayleigh.

References

- [1] Ahmad, A., Rather, A. A., Afful, R. B., Volodin, A. and Ali, I. (2024). Exploring the Fréchet–Power Rayleigh Distribution with Statistical Properties and Medical Data Insights. *Lobachevskii Journal of Mathematics*, 45(12), 6209-6223.
- [2] Marshall, A. W., Olkin, I. (1997). A new method for adding a parameter to a family of distributions with application to the exponential and Weibull families. *Biometrika*, 84(3), 641-652.

4th International Conference: Constructive Mathematical Analysis

Inference for the Reliability and Mean Time to Failure of Protected Consecutive k -out-of- n : G Systems

Erdem Cankut¹, Yunus Akdoğan¹, Kadir Karakaya¹ and Coşkun Kuş¹

¹ Department of Statistics, Faculty of Science, Selçuk University, Konya, Türkiye

erdem.cankut@selcuk.edu.tr, yakdogan@selcuk.edu.tr, kkarakaya@selcuk.edu.tr, coskun@selcuk.edu.tr

key-words: Consecutive k -out-of- n : G system, protection block, reliability.

Abstract:

This study develops inference for protected consecutive k -out-of- n :G systems under a complete lifetime sampling scheme. The known-parameter reliability structure is rewritten in terms of compact elementary blocks, which provide a convenient basis for estimation when the failure rates are unknown. By integrating these reliability blocks, compact closed-form expressions are obtained for the corresponding mean time to failure quantities. Using the resulting block representations, plug-in maximum likelihood estimators are constructed for the reliability and mean time to failure measures. The finite-sample behavior of these estimators is investigated by deriving their expected values and variances. An illustrative complete-data example demonstrates the implementation of the proposed estimators together with parametric bootstrap confidence intervals.

References

- [1] Casella, G., Berger, R. L. (2002). Statistical Inference (2nd ed.). Duxbury.
- [2] Eryilmaz, S. (2026). On reliability of consecutive k -out-of- n : G system equipped with protection blocks. International Journal of General Systems, 55(2), 129-144.
- [3] Kuş, C., Eryilmaz, S. (2026). Estimation of parameters for a system equipped with protection block. Journal of Computational and Applied Mathematics, 117102.

4th International Conference: Constructive Mathematical Analysis

A Study on Associated Curves and Their Curvature Relations

Aleyna Çetinkaya¹, Muhammed T. Sariaydin¹

¹ Department of Mathematics, Faculty of Science, Selçuk University, Konya, Türkiye

cetinkayaaleyna86@gmail.com, talatsariaydin@gmail.com

key-words: Associated curves, direction curves, Euclidean space.

Abstract:

This study, we introduces a class of associated curves constructed as integral trajectories of vector fields defined by the Positional Adapted Frame in Euclidean 3-space. The relationships between the curvature functions of these curves are examined in detail. Furthermore, the presented method provides constructive techniques for generating some special curves. Several illustrative examples are also presented to support the theoretical results.

References

- [1] Kandilci, G. T., Ekici, C. and Dede, M. (2023). Directional Bertrand Curves in Minkowski Space. Innovative Research in Natural Science and Mathematics, 15.
- [2] Korpınar, T., Sariaydin, M. T. and Turhan, E. (2013). Associated curves according to Bishop frame in Euclidean 3-space. Advanced Modeling and Optimization, 15(3), 713-717.
- [3] Sahiner, B. (2019). Direction curves of tangent indicatrix of a curve. Applied Mathematics and Computation, 343, 273-284.
- [4] Ünal, D., Ünlütürk, Y. and Ekici, C. (2025). A New Description of Q-Helices in Euclidean 4-Space. Differential Equations and Dynamical Systems, 1-15.

4th International Conference: Constructive Mathematical Analysis

Stability to Perturbations of Modified S -Fractions with Applications

Volodymyr Hladun¹, Marta Dmytryshyn² and Oleksandr Chaban¹

¹ Department of Applied Mathematics, Lviv Polytechnic National University, Lviv, Ukraine

² Department of Management and Administration, West Ukrainian National University, Ternopil, Ukraine

volodymyr.r.hladun@lpnu.ua, m.dmytryshyn@wunu.edu.ua, oleksandr.m.chaban@lpnu.ua

key-words: Modified S -fraction, stability to perturbations, set of uniform stability, relative error.

Abstract:

This presentation is devoted to the study of stability to perturbations for modified S -fractions (modified Stieltjes continued fractions) of the form

$$\frac{a_0}{z + \frac{a_1}{z + \frac{a_2}{z + \ddots}}}, \quad a_k > 0, \quad k \geq 0, \quad z \in \mathbb{C}. \quad (1)$$

While this class of functional continued fractions is a powerful tool in applied problems [1], its practical application requires a thorough understanding of the stability to perturbations [2].

We establish a formula quantifying the relative error of the approximants resulting from perturbations applied to the coefficients a_k of the modified S -fraction (1) itself. Utilizing this formula and leveraging techniques from the analytical theory of continued fractions, we derive sufficient conditions for the stability of (1) to perturbations.

We distinguish between two notions: the set of pointwise stability and the set of uniform stability. Specifically, assuming

$$0 < a_k \leq A, \quad 0 < \hat{a}_k \leq A, \quad k \geq 0,$$

where $\hat{a}_k$ are the perturbed coefficients, we prove that the set

$$\Omega_A = \{z \in \mathbb{C} : |\operatorname{Re} z| > \sqrt{A(1 - |\operatorname{Re} z|/|z|)}\}$$

is a set of stability to perturbations, and the set

$$\Omega_{\alpha, C} = \{z \in \mathbb{C} : |\operatorname{Im} z| \leq |\operatorname{Re} z| \tan \alpha, |\operatorname{Re} z| \geq C\}, \quad 0 \leq \alpha < \pi/2, \quad C > \sqrt{A(1 - \cos \alpha)},$$

is a set of uniform stability to perturbations of the modified S -fraction (1).

A significant outcome of this research is the derivation of explicit bounds for the relative errors of the approximants, which demonstrate how the perturbation error of a continued fraction's approximant depends on the magnitude of the perturbations and the values of the fraction's coefficients.

References

- [1] Cuyt, A., Petersen, V. B., Verdonk, B., Waadeland, H. and Jones, W. B. (2008). Handbook of Continued Fractions for Special Functions. Springer.
- [2] Dmytryshyn, M., Hladun, V. (2025). On the Sets of Stability to Perturbations of Some Continued Fraction with Applications. Symmetry, 17(9), 1442.

4th International Conference: Constructive Mathematical Analysis

A Chemotaxis-Navier-Stokes System with Dynamical Boundary Conditions

Baili Chen

Department of Mathematics, Computer Science and Statistics, Gustavus Adolphus College, Saint Peter, MN, USA

bchen@gustavus.edu

key-words: Dynamical boundary conditions, chemotaxis, Navier-Stokes.

Abstract:

A chemotaxis-Navier-Stokes system is studied under dynamical boundary conditions in a bounded convex domain with smooth boundary. This models the interaction of populations of swimming bacteria with the surrounding fluid. The existence of a global weak solution is proved using multiple layers of approximations and Rothe's method for the time discretization.

References

- [1] Vaz, C. L. D. (2007). Rothe's Method for an Isothermal Phase-field Model of a Binary Alloy with Convection. *Mathematica Contemporânea.*, 32, 221-251.
- [2] Vrábel, V., Šlodička, M. (2013). Nonlinear Diffusion Problem with Dynamical Boundary Value. *J. Comput. Appl. Math.*, 246 , 94-103.
- [3] Nečas, J. (1974). Application of Rothe's Method to Abstract Parabolic Equations. *Czechoslovak Mathematical Journal*, 24 (3), 496-500.
- [4] Tuval, I., Cisneros, L., Dombrowski, C., Wolgemuth, C. W., Kessler, J.O. and Goldstein, R. E. (2005). Bacterial Swimming and Oxygen Transport Near Contact Lines. *Proc. Natl. Acad. Sci. USA* (ISSN 0027-8424), 102 (7) 2277-2282.
- [5] Miranville, A., Zelik, S. (2005). Exponential Attractors for the Cahn-Hilliard Equation with Dynamic Boundary Conditions. *Math. Meth. Appl. Sci.*, 28 , 709-735.

4th International Conference: Constructive Mathematical Analysis

Korovkin-Type Approximation for Non-Positive Operators

Vijayalakshmi Balakrishnan Kiran Kumar¹, Memana Neelakantan Narayanan Namboodiri² and
Vinaya Poyil Chittarickal¹

¹ Department of Mathematics, Cochin University of Science and Technology, Cochin, India

² IMRT, Tiruvananthapuram and Formely of Cochin University of Science and Technology, Cochin, India

vbk@cusat.ac.in, mnnadri@gmail.com, vinayapc01@gmail.com

key-words: Korovkin-type theorems, non-positive operators, operator version, Grünwald operators.

Abstract:

The classical Korovkin theorem traditionally relies on the positivity of the underlying sequence of operators. However, in 1968, D. E. Wulbert established the first non-positive version. In this talk, I will present a generalized form of Wulbert's result to the class of uniformly bounded sequence of operators. As an application, I will discuss an operator version of this Korovkin-type theorem which will cover existing results in this direction. I will also present illustrative examples, which has its roots in the Grünwald's interpolation operator.

References

- [1] Faber, G. (1914). Über die interpolatorische Darstellung stetiger Funktionen. Jber. Deutsch. Math. Verein 23, 192–210.
- [2] Folland, G.B. (1999). Real analysis. Second edition, John Wiley & Sons, Inc., New York.
- [3] Grünwald, G. (1941). On a convergence theorem for the Lagrange interpolation polynomials. Bull. Amer. Math. Soc., 47, 271–275.
- [4] Grünwald, G. (1936). Über Divergenzerscheinungen der Lagrangeschen Interpolations polynome stetiger Funktionen. Ann. of Math. (2), 37, 908–918.
- [5] Kiran Kumar, V. B., Namboodiri, M. N. N. and Vinaya, P. C. (2024). Korovkin-type approximation for non-positive operators. arXiv:2403.03476
- [6] Korovkin, P. P. (1960). Linear operators and approximation theory. Russian Monographs and Texts on Advanced Mathematics and Physics, Vol. III, Gordon and Breach Publishers, New York.
- [7] Marcinkiewicz, J. (1937). Sur la divergence des polynomes d'interpolation. AM Kir. Ferencz József.
- [8] Phelps, R. R. (2001). Lectures on Choquet's Theorem. Lecture Notes in Mathematics, Vol. 1757, Springer-Verlag, Berlin, 2nd edition.
- [9] Wulbert, D. E. (1968). Convergence of operators and Korovkin's theorem. J. Approx. Theory, 1, 381–390.

4th International Conference: Constructive Mathematical Analysis

A Generalized Fourier-Feynman Transform over Paths in Abstract Wiener Space

Dong Hyun Cho

Department of Mathematics, Kyonggi University, Suwon, South Korea

j94385@kyonggi.ac.kr

key-words: Abstract Wiener space, analogue of Wiener measure, Fourier-Feynman transform, rotation theorem.

Abstract:

Let $(\mathbb{B}, \mathcal{B}(\mathbb{B}), \mu)$ be an abstract Wiener space and φ_a be a finite positive measure on $\mathcal{B}(\mathbb{B})$. Let $C^{\mathbb{B}}[a, b]$ denote an analogue of Wiener space over paths in abstract Wiener space, the space of $\mathbb{B}$ -valued continuous functions on $[a, b]$ and let $w_{a,b;\varphi_a}^{\rho,\beta,\mathbb{B}}$ be a generalized analogue of Wiener measure over $C^{\mathbb{B}}[a, b]$ according to β , φ_a and ρ , where $\rho > 0$ and β is a strictly increasing function on $[a, b]$. In this talk, we provide a rotation theorem on the product space $(C^{\mathbb{B}}[a, b])^2$ given by

$$\begin{aligned} & \int_{(C^{\mathbb{B}}[a,b])^2} F(\cos \theta y_1 - \sin \theta y_2, \sin \theta y_1 + \cos \theta y_2) d(w_{a,b;\varphi_a}^{\rho,\beta,\mathbb{B}})^2(y_1, y_2) \\ &= \int_{(C^{\mathbb{B}}[a,b])^2} F(y_1, y_2) d(w_{a,b;\varphi_a}^{\rho,\beta,\mathbb{B}})^2(y_1, y_2). \end{aligned}$$

where F is any (Borel) measurable function on $(C^{\mathbb{B}}[a, b])^2$. We then establish several results related to the equation just stated. This rotation theorem generalizes Bearman's rotation theorem and is essential to define the Fourier-Feynman transform on $C^{\mathbb{B}}[a, b]$ which plays a significant role in the study of Feynman integration theory, especially, in the study of probabilistic approach to Brownian motion.

References

- [1] Bearman, J. E. (1952). Rotation in the product of two Wiener spaces. Proc. Amer. Math. Soc., 3, 129–137.
- [2] Kuo, H. H. (1975). Gaussian measures in Banach spaces. Lecture Notes in Mathematics, Springer-Verlag, Berlin-New York, Vol. 463.
- [3] Ryu, K. S. (1992). The Wiener integral over paths in abstract Wiener space. J. Korean Math. Soc., 29(2), 317–331.
- [4] Ryu, K. S. (2010). Integration with respect to analogue of Wiener measure over paths in abstract Wiener space and its applications. Bull. Korean Math. Soc., 47(1), 131–149.

4th International Conference: Constructive Mathematical Analysis

A Neurodynamic Optimization Model for Smooth Pinball Loss Based Support Vector Machines

Vipavee Dammines¹, Pham Thi Vui² and Rabian Wangkeeree³

¹ Faculty of Science at Sriracha, Kasetsart University, Sriracha Campus, Chonburi, Thailand

² Department of Mathematics and Informatics, School of Education, Can Tho University, Can Tho, Vietnam

³ Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, Thailand

vipavee.d@ku.th, ptvui@ctu.edu.vn, rabianw@nu.ac.th

key-words: Pinball loss, smooth function, neurodynamic optimization.

Abstract:

Pinball loss function is widely used in Support Vector Machine (SVM) models due to its robustness; however, its non-smooth property limits the applicability of gradient-based optimization methods. In this research, a smooth approximation of pinball loss is developed and incorporated into the SVM framework to improve differentiability and optimization efficiency. Furthermore, a neurodynamic-based optimization approach is proposed to obtain the optimal solution through a stable dynamic system under Karush-Kuhn-Tucker (KKT) conditions. The proposed method is theoretically analyzed and experimentally evaluated using benchmark datasets from the UCI benchmark dataset. Numerical results demonstrate that the proposed model achieves superior convergence behavior and classification performance compared with existing state-of-the-art methods.

References

- [1] Nazemi, A., Dehghan, M. (2015). A neural network method for solving support vector classification problems. *Neurocomputing*, 152, 369-376.
- [2] Wangkeeree, R., Wangkeeree, R., Belay, Y. A., Ungchittrakool, K. and Nazemi, A. (2026). Neurodynamical neural network method for solving support vector machines with epigraph reformulated generalized pinball loss function. *Journal of Computational and Applied Mathematics*, 117754.
- [3] Xia, Y., Liu, Q., Wang, J. and Cichocki, A. (2024). A survey of neurodynamic optimization. *IEEE Transactions on Emerging Topics in Computational Intelligence*, 8(4), 2677-2696.

4th International Conference: Constructive Mathematical Analysis

Automatic Differentiation Using Neural Networks and Their Comparison with Higher Order Spectral Differentiation

Mrinal Kanti Das¹, Lokendra Kumar Balyan¹

¹ Discipline of Natural Science (Mathematics), PDPM Indian Institute of Information Technology, Design and Manufacturing, Jabalpur, India

23pnme02@iiitdmj.ac.in, balyan@iiitdmj.ac.in.

key-words: Automatic differentiation, back propagation, spectral differentiation.

Abstract:

Automatic differentiation (AD), also called algorithmic differentiation, applies the chain rule from calculus step by step to compute the derivative while the program is running. That is why it performs faster and more accurately compared to other differentiations and is also easy to use. Calculate the derivative value and the function value simultaneously. AD is widely used in Physics-Informed Neural Networks (PINNs) and has important applications in computational fluid dynamics, atmospheric sciences, and engineering design optimisation. In this paper, we consider two continuous and two discontinuous functions. We will apply both the spectral derivative and AD methods to those functions and compare the results graphically. The spectral derivative matches very closely with AD for smooth functions. AD results are much better compared to the classical method, which shows oscillations near the boundary and at discontinuous points.

References

- [1] Baydin, A. G., Pearlmutter, B. A., Radul, A. A. and Siskind, J. M. (2018). Automatic Differentiation in Machine Learning: A Survey. *Journal of Machine Learning Research*, 18(153), 1–43.
- [2] Boyd, J. P. (2000). *Chebyshev and Fourier Spectral Methods* (2nd ed.). Dover Publications.
- [3] Paszke, A., Gross, S., Chintala, S., Chanan, G., Yang, E., DeVito, Z., Lin, Z., Desmaison, A., Antiga, L. and Lerer, A. (2017). Automatic Differentiation in PyTorch. *NIPS 2017 Autodiff Workshop: The Future of Gradient-Based Machine Learning Software and Techniques*.
- [4] Raissi, M., Perdikaris, P. and Karniadakis, G. E. (2019). Physics-Informed Neural Networks: A Deep Learning Framework for Solving Forward and Inverse Problems Involving Nonlinear Partial Differential Equations. *Journal of Computational Physics*, 378, 686–707.
- [5] Trefethen, L. N. (2000). *Spectral Methods in MATLAB*. Society for Industrial and Applied Mathematics (SIAM).

4th International Conference: Constructive Mathematical Analysis

Fixed-Point Iterations for Analyzing the Topology of Biological Networks

Tugce Delen¹, Ozlem Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

tugcedelen@yahoo.com, acarozlem@gmail.com

key-words: Fixed point theory, mathematical modeling, network topology.

Abstract:

Cellular decisions and metabolic balances emerge from the collective, rhythmic, and nonlinear interactions of numerous chemical components. Biological networks model this complexity as directed graphs $G = (V, E)$, where vertices represent intracellular components (proteins, enzymes, metabolites) and edges represent activation or inhibition relationships. The insulin signaling pathway in Type-2 Diabetes is a high-dimensional, nonlinear regulatory network governed by nested feedback loops.

The dynamic behavior of this system is modeled by nonlinear ordinary differential equations (ODEs):

$$\frac{dx_i}{dt} = f_i(x_1, x_2, \dots, x_n) - g_i(x_i), \quad i = 1, \dots, n,$$

where $x_i(t) \in \mathbb{R}_{\geq 0}$ denotes concentrations, f_i is a nonlinear Hill-type production function, and g_i represents the degradation rate. The fixed points $\mathbf{x}^* = (x_1^*, x_2^*, \dots, x_n^*)^T$ satisfy $\frac{dx_i}{dt} = 0$. A stable fixed point corresponds to homeostasis, whereas an unstable fixed point or bifurcation signals pathogenesis. However, analytically determining fixed points in high-dimensional systems is difficult due to the complex Jacobian matrix $J \in \mathbb{R}^{n \times n}$.

To overcome this, we apply the Circuit-Breaking Algorithm (CBA) (Radde 2010, 2012), bridging dynamical systems and graph theory. Here, we adapt this framework to insulin signaling pathways to analytically identify critical bifurcation thresholds triggering pathogenesis, offering a scalable tool for complex human disease network analysis [1, 2, 3]. **Acknowledgements:** The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Kummer, U., Olsen, L. F., Dixon, C. J., Green, A. K., Bornberg-Bauer, E. and Baier, G. (2000). Switching from simple to complex oscillations in calcium signaling. *Biophysical Journal*, 79(3), 1188–1195.
- [2] Radde, N. (2010). Fixed point characterization of biological networks with complex graph topology. *Bioinformatics*, 26(22), 2874–2880.
- [3] Radde, N. (2012). Analyzing fixed points of intracellular regulation networks with interrelated feedback topology. *BMC Systems Biology*, 6(1), 57.

4th International Conference: Constructive Mathematical Analysis

New Results on Fixed Point Theory in Perturbed Metric Spaces with Directed Graphs

Tugce Delen¹, Ozlem Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

tugcedelen@yahoo.com, acarozlem@ymail.com

key-words: Perturbed metric spaces, fixed point theory, directed graphs.

Abstract:

This paper establishes new fixed point results for Geraghty-type contractive mappings on complete perturbed metric spaces endowed with directed graph structures. By integrating the concepts of perturbed metrics and graph-preserving mappings, a generalized Banach–Geraghty fixed point theorem is obtained, ensuring the existence and uniqueness of fixed points under suitable contractive conditions. The proposed framework simultaneously captures perturbation effects in distance measurements and structural relationships represented by directed graphs, thereby extending several classical fixed point results in metric and graph-based settings. Furthermore, the applicability of the theoretical findings is demonstrated through a nonlinear Fredholm integral equation, illustrating the effectiveness of the developed approach in the analysis of nonlinear problems involving uncertainty and interconnected structures.

References

- [1] Acar, Ö., Aydi, H. and De la Sen, M. (2021). New fixed point results via a graph structure. *Mathematics*, 9(9), 1013.
- [2] Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3, 133–181.
- [3] Geraghty, M. (1973). On contractive mappings. *Proceedings of the American Mathematical Society*, 40(2), 604–608.
- [4] Jachymski, J. (2008). The contraction principle for mappings on a metric space with a graph. *Proceedings of the American Mathematical Society*, 136, 1359–1373.
- [5] Jleli, M., Samet, B. (2025). On Banach’s fixed point theorem in perturbed metric spaces. *Journal of Applied Analysis and Computation*, 15, 993–1001.
- [6] Păcurar, C., Târnoveanu, M. A. (2025). Mappings contracting perimeters of triangles in perturbed metric spaces. *Altay Conference Proceedings in Mathematics*, 1(1), 68–72.

4th International Conference: Constructive Mathematical Analysis

On the Approximations of the Ratios of Double Hypergeometric Functions by Branched Continued Fractions

Roman Dmytryshyn

Department of Mathematical and Functional Analysis, Vasyl Stefanyk Carpathian National University, Ivano-Frankivsk, Ukraine

roman.dmytryshyn@cnu.edu.ua

key-words: Double hypergeometric functions, branched continued fractions, approximation by rational functions, convergence, numerical stability.

Abstract:

The problems of approximating the ratios of double hypergeometric series by branched continued fractions are considered: construction of the simplest fractional structure and investigation of the convergence and numerical stability of the constructed expansion. New methods are proposed for investigating the convergence and establishing the sets of numerical stability of the expansions of the ratios of some double hypergeometric series, including the well-known Appell hypergeometric series, into branched continued fractions [1].

References

- [1] Dmytryshyn, R. (2025). Truncation error bounds for branched continued fraction expansions of some Appell's hypergeometric functions F_2 . Symmetry, 17(8), 1204.

4th International Conference: Constructive Mathematical Analysis

Solution Structure of Higher-Order Nonlinear Hadamard-Type Fractional p -Laplacian Problems on Finite Domains

Mürüvvet Tuva Erbaş^{1,2}, Erbil Çetin¹

¹ Department of Mathematics, Ege University, İzmir, Türkiye

² Department of Mathematics, Hatay Mustafa Kemal Üniversitesi, Hatay, Türkiye

muruvvettuva.eras@mku.edu.tr, erbil.cetin@ege.edu.tr

key-words: Hadamard fractional derivative, nonlinear p -laplacian, fixed point methods.

Abstract:

We investigate a class of higher-order nonlinear boundary value problems involving Hadamard-type fractional operators and the p -Laplacian on a finite interval subject to integral boundary conditions. The problem is reformulated as an equivalent integral equation and analyzed within an operator-theoretic framework in a suitable Banach space.

Two independent solvability results are established. The first guarantees existence and uniqueness of solutions under standard continuity and growth assumptions via a contraction principle. Unlike earlier studies relying solely on Banach contraction arguments, the present analysis establishes existence results for fully nonlinear configurations under weaker hypotheses. The second result therefore provides the existence of at least one solution under more general nonlinear structures through alternative operator estimates.

This framework enables the treatment of fully nonlinear higher-order problems together with integral-type boundary conditions, extending several earlier results restricted to semilinear or lower-order settings. An illustrative example is provided to demonstrate the applicability of the theoretical findings. The obtained results contribute to the constructive theory of Hadamard-type fractional differential equations and provide a rigorous analytical foundation for nonlinear fractional p -Laplacian models.

Acknowledgment. The first author would like to thank TUBITAK for supporting her as a scholar during her PhD through the 2211-E Domestic Direct Doctoral Fellowship Program.

References

- [1] Pei, K., Wang, G. and Sun, Y. (2017). Successive iterations and positive extremal solutions for a Hadamard-type fractional integro-differential equations on infinite domain. *Applied Mathematics and Computation*, 312, 158–168.
- [2] Şenlik Çerdik, T., Yörük Deren, F. (2022). New results for higher-order Hadamard-type fractional differential equations on the half-line. *Mathematical Methods in the Applied Science*, 45(4), 2315–2330.
- [3] Thiramanus, P., Ntouyas, S. K. and Tariboon, J. (2016). Positive solutions for Hadamard fractional differential equations on infinite domain. *Advances in Difference Equations*, 2016:83, 1–18.
- [4] Xu, J., Cui, Y. and O'Regan, D. (2025). Positive solutions for a Hadamard-type fractional-order three-point boundary value problem on the half-line. *Nonlinear Analysis: Modelling and Control*, 30(2), 176–195.
- [5] Zhang, W., Liu, W. (2018). Existence of solutions for several higher-order Hadamard-type fractional differential equations with integral boundary conditions on infinite interval. *Boundary Value Problems*, 2018:134, 1–27.

4th International Conference: Constructive Mathematical Analysis

XBilal Distribution: Properties, Parameter Estimation and Applications

Tenzile Erbayram¹, Yunus Akdogan¹

¹ Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

tenzile.erbayram@selcuk.edu.tr, yakdogan@selcuk.edu.tr

key-words: Estimation, finite mixture distribution, Monte Carlo simulation.

Abstract:

This study presents the XBilal distribution, which is obtained as a special finite mixture of exponential and Bilal distributions. The fundamental mathematical and statistical properties of this new distribution, such as the probability density, moment generating, survival, and hazard function, have been derived. Different point estimators, including the maximum likelihood, least squares, weighted least squares, Cramér-von Mises, Anderson-Darling, and maximum product of spacings have been considered for estimating the model parameters. To empirically support the theoretical findings of the proposed distribution, a comprehensive Monte Carlo simulation study is conducted. This simulation process is implemented under different sample sizes and parameter combinations, comparing the performance of the estimators considered. The efficiency, consistency, and relative superiority of the estimators are analyzed using statistical performance criteria such as bias and mean squared errors. Finally, an application study is carried out on a real dataset to demonstrate the flexibility of the proposed distribution and its superiority over traditional models. The findings show that the XBilal distribution provides a better fit than existing models and is an effective alternative in data modeling.

References

- [1] Everitt, B. S. (1996). An introduction to finite mixture distributions. Statistical methods in medical research, 5(2), 107-127.
- [2] McLachlan, G. J., Peel, D. (2000). Finite mixture models. New York, NY: John Wiley & Sons.

4th International Conference: Constructive Mathematical Analysis

On a Statistical Versions of Certain Types of Alpha Convergence

Alper Erdem¹, Tuncay Tunç¹

¹ Department of Mathematics, Mersin University, Mersin, Türkiye

alpererdem@mersin.edu.tr, ttunc@mersin.edu.tr

key-words: Alpha convergence, function sequences, statistical convergence.

Abstract:

This study considers new types of alpha type convergences in the context of statistical convergence for sequences of functions defined between metric spaces. Furthermore, using this concepts, we prove some fundamental properties and get connections between them.

Let (X, d) and (Y, ρ) be metric spaces, (f_n) be a sequence of functions and f be a function from X to Y .

Definition 1. [1] The function sequence (f_n) is called *uniform alpha convergent* to the function f on X if for every sequence (x_n) and (y_n) of points of X with the property $d(x_n, y_n) \rightarrow 0$, the sequence $\rho(f_n(x_n), f(y_n)) \rightarrow 0$.

Definition 2. [1] The function sequence (f_n) is called *Cauchy alpha convergent* to the function f on X if for every Cauchy sequence (x_n) and (y_n) of points of X with the property $d(x_n, y_n) \rightarrow 0$, the sequence $\rho(f_n(x_n), f(y_n)) \rightarrow 0$.

Definition 3. [2] The sequence $(f_n) \in \mathfrak{sf}(Y^X)$ is called *semi uniform alpha convergent* to $f \in Y^X$ on X and denoted by $f_n \rightrightarrows_{s\alpha} f$ if

1. $f_n \rightarrow f$
2. For every $\varepsilon > 0$ there exists a $\delta = \delta(\varepsilon) > 0$ such that for all $n \in \mathbb{N}$ there exists $m \in \mathbb{N}$ with $m \geq n$ such that for all $x, y \in X$ with $d(x, y) < \delta$ we have $\rho(f_m(x), f(y)) < \varepsilon$

hold.

References

- [1] Erdem, A., Tunç, T. (2025). On Cauchy alpha and uniform alpha convergence. Filomat, 39(20), 7033–7042.
- [2] Erdem, A., Arpacioğlu, A., and Tunç, T. (2026). On Semi Uniform Alpha Convergence and a Korovkin-Type Theorem. Dolomites Research Notes on Approximation, 19(2), 44–51.

4th International Conference: Constructive Mathematical Analysis

Convergence and Asymptotic Behavior of Mellin-Steklov Exponential Sampling Series in Logarithmically Weighted Spaces

Dilek Erdogan¹, Tuncer Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

dilekozer@yahoo.com, tunceracar@ymail.com

key-words: Mellin-Steklov integrals, exponential sampling series, weighted approximation.

Abstract:

In this talk, we investigate the approximation properties of r -th order ($r \in \mathbb{N}$) Mellin-Steklov type exponential sampling operators within logarithmically weighted spaces of continuous functions. Initially, we establish the well-definiteness of these operators, demonstrate their pointwise and uniform convergence in weighted spaces, and present the rates of convergence via a suitable modulus of continuity in logarithmic weighted spaces. Subsequently, we establish a quantitative representation of the pointwise asymptotic behavior of these operators using Mellin-Taylor's expansion.

References

- [1] Acar T., Kursun S. (2023). Pointwise convergence of generalized Kantorovich exponential sampling series. Dolomites Research Notes on Approximation, 16(2), 1-10.
- [2] Acar, T., Kursun, S., and Acar, Ö. (2024). Approximation properties of exponential sampling series in logarithmic weighted spaces. Bulletin of the Iranian Mathematical Society, 50(3), 36.
- [3] Angamuthu, S.K., Bajpeyi, S. (2020). Direct and inverse result for Kantorovich type exponential sampling series. Results Math., 75(3).
- [4] Aral, A., Acar, T. and Kursun, S. (2022). Generalized Kantorovich forms of exponential sampling series. Anal.Math.Phys., 12(2).
- [5] Bardaro, C., Faina, L. and Mantellini I. (2017) A generalization of the exponential sampling series and its approximation properties, Math. Slovaca, 67(6), 1481-1496.
- [6] Costarelli, D. (2024) Convergence and high order of approximation by Steklov sampling operators. Banach J. Math. Anal., 18:70.
- [7] Erdogan, D., Kursun, S. and Acar, T. (2026) Approximation properties of Mellin-Steklov type exponential sampling series. Submitted.

4th International Conference: Constructive Mathematical Analysis

Approximation of Discontinuous Functions By Kantorovich Exponential Neural Network Operators

Dilek Erdogan¹, Serife Nur Karaman¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

dilekozer@yahoo.com, serifenurkaraman@yahoo.com

key-words: Kantorovich exponential neural networks operators, jump discontinuity, sigmoidal function.

Abstract:

In this talk, we analyze the behavior of Kantorovich exponential neural network operators when discontinuous signals are considered. First, we establish a fundamental lemma that decomposes the operator by utilizing the critical index at the point of a jump discontinuity. Using this lemma, we prove certain approximation theorems, providing necessary and sufficient conditions for the convergence of these operators at non-removable jump discontinuities. Furthermore, we investigate the conditions under which the approximation is not achieved; specifically, we present a theorem demonstrating that the family of operators fails to converge pointwise at the discontinuity point without strict additional assumptions on the sigmoidal kernel. Finally, some numerical computations are performed to verify the approximation of discontinuous functions f by the proposed operators $K_n^{\chi\sigma} f$.

References

- [1] Agrawal, P. N., Baxhaku, B. and Berisha, A. (2025). Enhanced function approximation and applications to image scaling: A new family of exponential sampling neural network Kantorovich operators. *Applied Soft Computing*, 174, 112923.
- [2] Kumar, A. S., Kumar, P. and Devaraj, P. (2022). Approximation of discontinuous functions by Kantorovich exponential sampling series: AS Kumar et al. *Analysis and Mathematical Physics*, 12(3), 73.
- [3] Kursun, S., Acar, T. (2025). Approximation of discontinuous signals by exponential-type generalized sampling Kantorovich series. *Mathematical Methods in the Applied Sciences*, 48(1), 340-355.

4th International Conference: Constructive Mathematical Analysis

Fixed Point Results for Rational Type F -Contractions in Perturbed Metric Spaces

Esref Erdogan¹, Ozlem Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

esreferdogan@yahoo.com, acarozlem@ymail.com

key-words: Fixed point, perturbed mapping, perturbed metric space.

Abstract:

This study investigates fixed point theory within the framework of perturbed metric spaces, a setting that extends classical metric spaces by incorporating systematic perturbations into distance measurements. We introduce a new class of mappings termed "rational type F -contractions," which generalizes Wardowski's F -contractions by utilizing a rational expression in the contraction condition. The main result establishes the existence and uniqueness of fixed points for these mappings in complete perturbed metric spaces, provided the mapping is perturbed continuous. Furthermore, we present illustrative examples and corollaries to demonstrate that our findings unify and extend several existing fixed point theorems in the literature.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Banach, S. (1922). Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3, 133–181.
- [2] Wardowski, D. (2012). Fixed points of a new type of contractive mappings in complete metric spaces. *Fixed Point Theory and Applications*, 2012, 1-6.
- [3] Jleli, M., Samet, B. (2025). On Banach's fixed point theorem in perturbed metric spaces. *J. Appl. Anal. Comput.*, 15, 993-1001.

4th International Conference: Constructive Mathematical Analysis

Fixed Point Theorems for Z -Contractions in Perturbed Metric Spaces

Eşref Erdoğan¹, Aybala Sevd e Özkapu¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

esreferdogan@yahoo.com, aybalasevdeozkapu@yahoo.com

key-words: Fixed point, perturbed mapping, perturbed metric space.

Abstract:

In this presentation, we extend the concept of simulation functions and Z -contractions to the framework of perturbed metric spaces. Perturbed metric spaces, which generalize classical metric structures by incorporating a perturbation term into the standard triangle inequality, provide a more flexible and adaptable setting for fixed point theory. We introduce the notion of Z -contractions in perturbed metric spaces and establish new criteria for the existence and uniqueness of fixed points for such mappings. To demonstrate the validity and effectiveness of our theoretical results, we provide a concrete non-trivial example within a perturbed metric space where standard metric axioms are appropriately relaxed. Ultimately, our findings unify and generalize several existing fixed point theorems in the literature, effectively bridging the gap between simulation functions and perturbed metric structures.

References

- [1] Jleli, M., Samet, B. (2025). On Banach's fixed point theorem in perturbed metric spaces. *J. Appl. Anal. Comput.*, 15, 993-1001.
- [2] Khojasteh, F., Shukla, S., Radenović, S. (2015). A new approach to the study of fixed point theory for simulation functions, *Filomat*, 29(6), 1189–1194.

4th International Conference: Constructive Mathematical Analysis

An Affine ARX-Based Adaptive RST Framework for Disturbed Buck Converter Systems

Fatih Evran

Department of Electrical and Electronics Engineering, Düzce University, Düzce, Türkiye

fatih evran@duzce.edu.tr

key-words: Affine discrete-time systems, recursive identification, adaptive polynomial control.

Abstract:

This study considers an adaptive polynomial control problem for a nonideal DC–DC buck converter operating under line and load disturbances. Starting from a nonideal averaged continuous-time model, an equivalent affine discrete-time input–output representation is introduced for control design purposes. The proposed model is formulated as a bias-augmented ARX structure, where the additional constant term is used to capture steady-state offsets, operating-point shifts, and lumped nonideal effects that cannot be adequately represented by a low-order linear ARX model alone. This formulation allows a separation between dynamic coefficients and constant offset components, thereby reducing unnecessary parameter drift in online identification. Based on this representation, the converter dynamics are described by time-varying polynomial coefficients that are recursively estimated via a forgetting-factor recursive least-squares algorithm. The estimated coefficients are then used to update, at each sampling instant, the RST controller polynomials through a Diophantine equation associated with a prescribed closed-loop characteristic polynomial. The resulting controller includes an explicit discrete-time integral factor, which ensures offset rejection and asymptotic tracking of constant references under constant disturbances, provided that the closed-loop system remains stable. The proposed framework yields a unified adaptive control scheme in which the bias term improves the identification quality, while the polynomial controller compensates for operating-point variations induced by abrupt input-voltage and load changes. The analysis shows that the affine augmentation is not merely a modeling convenience, but a structural component that improves parameter interpretability and supports smoother online adaptation. The effectiveness of the proposed scheme is evaluated through simulation studies under line and load disturbances.

References

- [1] Abdurraqueeb, A. M., Al-Shamma'a, A. A., Alkuhayli, A., Noman, A. M. and Addoweesh, K. E. (2022). RST digital robust control for DC/DC buck converter feeding constant power load. *Mathematics*, 10(10), 1782.
- [2] Das Neves, C. C., Junior, W. B., de Medeiros, R. L. P., Ayres Junior, F. A. C., Bessa, I. V., Veroneze, G. de M., Silva, L. E. S. and Farias, N. J. S. (2020). Direct form digital robust RST control based on Chebyshev sphere optimization applied in a DC-DC power converter. *Energies*, 13(15), 3810.
- [3] Hajizadeh, A., Shahirinia, A. H., Namjoo, N. and Yu, D. C. (2015). Self-tuning indirect adaptive control of non-inverting buck–boost converter. *IET Power Electronics*, 8(11), 2299–2306.
- [4] Kermadi, M., Rebai, A., Baghli, L., Mesbahi, N., Mekhilef, S., Mubin, M. and Mokhlis, H. (2024). Model-free predictive current controller for voltage source inverters using ARX model and recursive least squares. *IEEE Transactions on Circuits and Systems II: Express Briefs*, 71(5), 2619–2623.

- [5] Kermadi, M., Rebai, L., Baghli, L., Mesbahi, N., Mekhilef, S., Mubin, M., Mokhlis, H. and Rondan, H. S. R. (2025). Model-free deadbeat predictive current control for grid-connected inverters using autoregressive model and recursive least squares. *IEEE Journal of Emerging and Selected Topics in Power Electronics*, 13(4), 5138–5148.
- [6] Rodriguez, J., Heydari, R., Rafiee, Z., Young, H. A., Flores-Bahamonde, F. and Shahparasti, M. (2020). Model-free predictive current control of a voltage source inverter. *IEEE Access*, 8, 211104–211114.

4th International Conference: Constructive Mathematical Analysis

Dominants and Their Properties in Lattice Scaled Spaces

Niyazi Anıl Gezer

Department of Mathematics, Faculty of Arts and Sciences, TED University, Ankara, Türkiye.

anilgezer@gmail.com

key-words: Dominated Map, ultrametric space, discrete dynamics.

Abstract:

In the present talk, we focus on L -scaled spaces, a class of generalized ultrametric spaces where the distance function $d : M \times M \rightarrow L$ takes values in a lattice L possessing a minimal element, see [1]. These spaces provide a robust setting for analyzing the bo -convergence of sequences, which lifts the classical σ -convergence on L to M . The primary focus of this talk is the theory of domination, wherein the behavior of a mapping $f : M \rightarrow M$ is governed by an order-theoretic modulus of continuity known as a dominant (or majorant) $S : L \rightarrow L$. We establish that for a specific mapping, the collection of all dominants inherits the lattice structure of L and is closed under pointwise order limits. This duality between the mapping's dynamics and the dominants' algebraic properties forms the basis for characterizing σ - bo -continuity and the preservation of bo -Cauchy sequences. Furthermore, we present several metric inequalities in L that do not hold in classical metric settings.

References

- [1] Gezer, N. A. (2025). Majorants, Uniformities, and bo -Convergence in L -Scaled Spaces. Turkish Journal of Mathematics and Computer Science, 17(2), 304-309.

4th International Conference: Constructive Mathematical Analysis

Numerical Stability of Branched Continued Fraction Expansions of Lauricella-Saran Hypergeometric Functions F_K and Their Ratios in Special Cases

Vitaliy Goran

Department of Mathematical and Functional Analysis, Vasyl Stefanyk Carpathian National University, Ivano-Frankivsk, Ukraine

vitaliy.goran.23@pnu.edu.ua

key-words: Numerical stability, backward recurrence algorithm, branched continued fraction, Lauricella-Saran's hypergeometric function F_K , approximation by rational functions.

Abstract:

The numerical stability of the backward recurrence algorithm for computing approximants expansions of the Lauricella-Saran hypergeometric functions F_K and their ratios in special cases into branched continued fractions is considered. New estimates of the relative errors of approximants computations are obtained, showing the dependence of the error of the approximant on the magnitude of the rounding errors of its elements, and the values of the coefficients of the branched continued fractions. New sufficient conditions for the sets of numerical stability for the aforementioned approximants are also established. Numerical experiments allowed us to evaluate the practical effectiveness of the proposed theoretical results.

References

- [1] Dmytryshyn, R. Goran, V. (2026) Numerical stability of branched continued fraction expansions of Lauricella-Saran's hypergeometric function F_K ratios. Demonstr. Math., 59(1), 20250224.

4th International Conference: Constructive Mathematical Analysis

Some Properties of Generalized Degenerate Tangent Polynomials and Their Associated Matrices

Juan Hernández¹, Yamilet Quintana² and William Ramírez³

¹ Escuela de Matemáticas, Universidad Autónoma de Santo Domingo, Santo Domingo, Dominican Republic

² Department of Mathematics, Universidad Carlos III de Madrid, Madrid, Spain

³ Department of Natural and Exact Sciences, Universidad de la Costa, Barranquilla, Colombia

jhernandez14@uasd.edu.do, yaquinta@math.uc3m.es, wramirez4@cuc.edu.co

key-words: Generalized degenerate tangent polynomials, tangent polynomials, ∇_λ -Appell sequences.

Abstract:

We introduce and study generalized degenerate tangent polynomials $T_{s,\lambda}^{(\alpha)}(\xi)$ and their associated matrices using the generating function approach. Using this framework, we derive their main algebraic and differential properties, including their characterization as Δ_λ -Appell sequences, recurrence relations, inversion formulas, and addition formulas. We also obtain new factorizations and matrix identities connecting generalized degenerate Pascal matrices, tangent polynomial matrices, and Fibonacci matrices. Numerical experiments further illustrate the symmetry and distribution of their zeros in the complex plane.

References

- [1] Cesarano, C., Quintana, Y. and Ramírez, W. (2024). Degenerate versions of hypergeometric Bernoulli–Euler polynomials. *Lobachevskii J. Math.*, 45(8), 3509–3521.
- [2] Hernández, J., Peralta, D. and Quintana, Y. (2023). A look at generalized degenerate Bernoulli and Euler matrices. *Mathematics*, 11(12), 2731.
- [3] Hernández, J., Quintana, Y. and Ramírez, F.J. (2024). A note on generalized degenerate q -Bernoulli and q -Euler matrices. *Dolomites Res. Notes Approx.* 17, 63–71.
- [4] Hernández, J., Quintana, Y. and Ramírez, W. (2026). Generalized degenerate tangent polynomials and their associated matrices. Submitted.
- [5] Ryoo, C. S. (2017). Appell-type degenerate tangent polynomials. *Appl. Math. Sci.*, 11(54), 2671–2679.
- [6] Ryoo, C. S. (2015). Notes on degenerate tangent polynomials. *Global J. Pure Appl. Math.*, 11(5), 3631–3637.
- [7] Simsek, Y. (2020). Some new families of special polynomials and numbers associated with finite operators. *Symmetry*, 12(2), 237.

International Conference: Constructive Mathematical Analysis

Structural Stability and Rational q -Deformations in Parikh q -Determinants

Seyda Ildan¹, Aynur Yalciner¹

¹ Department of Mathematics, Selçuk University, Konya, Türkiye

seydaildan@selcuk.edu.tr, ayalciner@selcuk.edu.tr

key-words: Parikh q -matrix, q -Stirling numbers, Hessenberg matrix, Euler operator.

Abstract:

In this study, we investigate the degree mismatch phenomenon arising in the classical q -Stirling expansions of Parikh q -determinants associated with periodic words. For determinant families generated by periodic structures of the form $w = (ba)^n$, the synchronization of the determinant sequence with q is introduced and its analytic extension modeled via the Euler differential operator $T = q \frac{d}{dq}$ is examined. Then, the connections between these analytic structures and weighted Hessenberg-type matrices are addressed and various structural stability constraints are derived in this context. In particular, the degree-suppressing effects of lexicographical inversions ($b \rightarrow a$) on the determinant behavior are investigated. Using a structural asymmetry index (τ), we propose structural stability criteria describing when periodic structures remain inside the polynomial ring $\mathbb{Z}[q]$ and when rational q -deformations arise in the field $\mathbb{Q}(q)$. These results establish new connections between Parikh q -determinants, operator methods, generating functions and structural stability phenomena in combinatorics on words.

References

- [1] Atanasiu, A., Martin-Vide, C. and Mateescu, A. (2001). On the injectivity of the Parikh matrix mapping. *Fundamenta Informaticae*, 49(3), 289-299.
- [2] Egecioğlu, Ö., Ibarra, O. H. (2005). The Parikh q -matrix of a word. *Advances in Applied Mathematics*, 34(3), 624-645.

4th International Conference: Constructive Mathematical Analysis

On the Zero Behavior of Degenerate Bernstein Basis Functions

Rodrigo Ingelmo¹, Yamilet Quintana¹

¹ Department of Mathematics, Universidad Carlos III de Madrid, Madrid, Spain

100516064@alumnos.uc3m.es, yaquinta@math.uc3m.es

key-words: Degenerate Bernstein basis functions, Bernstein basis functions, zeros and critical points.

Abstract:

The distribution of zeros of a polynomial plays a central role in characterizing its structural and analytic properties, particularly in relation to how these features evolve under parameter variations. This study focuses on deformations of Bernstein polynomials induced by a deformation parameter λ , interpolating between classical and generalized settings. Within this framework, we analyze the zero distribution of degenerate Bernstein basis functions in the sense of T. Kim and D. Kim [4, 5], together with the behavior of their associated critical points. We establish sharp localization results, showing that these zeros lie in explicitly described λ -dependent intervals. In addition, we prove a precise interlacing between zeros and critical points and characterize its evolution as λ varies. These results provide a quantitative description of the behavior of zeros and critical points with respect to the deformation parameter.

References

- [1] Allan, L. (2021). Approximation and interpolation with Bernstein polynomials. Baylor University, USA. Doctoral Dissertation.
- [2] Ingelmo, R. (2024). Algebraic and analytic properties of degenerate Bernstein polynomials. Universidad Carlos III de Madrid, Spain. Unpublished master's thesis.
- [3] Ingelmo, R., Quintana, Y. (2026). Degenerate Bernstein forms: some computational features and stochastic logic associated. Kragujevac J. Math. Online first.
- [4] Kim, T., Kim, D. (2019). Degenerate Bernstein polynomials. Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM, 113(3), 2913–2920.
- [5] Kim, T., Kim, D., Jang, G. and Kwon, J. (2019). A note on degenerate Bernstein polynomials. J. Inequal. Appl. 1, 1–12.

4th International Conference: Constructive Mathematical Analysis

Approximation Properties of Modified Neural Network Operators Activated by Sigmoidal Functions

Serife Nur Karaman¹, Tuncer Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

serifenurkaraman@yahoo.com, tunceracar@ymail.com

key-words: Neural network operators, sigmoidal functions, modulus of continuity.

Abstract:

In this study, we introduce and investigate a novel family of linear neural network operators activated by sigmoidal functions and modified by a suitable function ρ . This modification yields a non-uniform sampling scheme, enabling the approximation process to adapt effectively to the local behavior of the target function. We establish the fundamental approximation properties of these operators, proving both pointwise and uniform convergence on compact intervals. Furthermore, a quantitative error estimate is derived via the modulus of continuity, revealing that the rate of convergence is governed by the smoothness of the transformed function $f \circ \rho^{-1}$. We also establish a Voronovskaya-type asymptotic formula that explicitly characterizes the local approximation error in terms of the first and second derivatives of ρ . Finally, numerical examples are presented to validate the theoretical findings and demonstrate the efficacy of the proposed operators in selective approximation tasks.

References

- [1] Cárdenas-Morales, D., Garrancho, P. and Raşa, I. (2011). Bernstein-type operators which preserve polynomials. *Computers and Mathematics with Applications*, 62(1), 158-163.
- [2] Costarelli, D., Spigler, R. (2013). Approximation results for neural network operators activated by sigmoidal functions. *Neural Networks*, 44, 101-106.
- [3] Turgay, M., Acar, T. (2024). Approximation by modified generalized sampling series. *Mediterranean Journal of Mathematics*, 21(3), 107.

4th International Conference: Constructive Mathematical Analysis

Convergence of Multidimensional Mellin Type Singular Integral Operators with Non-Isotropic Kernels

Harun Karsli

Department of Mathematics, Bolu Abant İzzet Baysal University, Bolu, Türkiye

karsli_h@ibu.edu.tr

key-words: Linear singular integral, Non-isotropic distance, multidimensional Mellin convolution integral, convergence, rate of convergence.

Abstract:

Here we give some approximation theorems concerning pointwise convergence and rate of pointwise convergence of multidimensional Mellin type singular integral operators of the form

$$(T_w f)(\mathbf{s}) = \int_{\mathbb{R}_+^n} K_w(\mathbf{t}) f(\mathbf{s}\mathbf{t}) \frac{d\mathbf{t}}{\langle \mathbf{t} \rangle}, \quad w > 0,$$

where the kernel function K_w satisfies some singularity assumptions with non-isotropic kernels.

References

- [1] Acar, T., Kursun, S. and Acar, O. (2024) Approximation properties of exponential sampling series in logarithmic weighted spaces. Bull Iran Math Soc., 50:36.
- [2] Anastassiou, G. A., Aral, A. (2009). Generalized Picard singular integrals. Comput. Math. Appl., 57(5), 821–830.
- [3] Angeloni, L., Vinti, G. (2014). Variation and Approximation in Multidimensional Setting for Mellin Integral Operators. In: Zayed, A., Schmeisser, G. (eds), New Perspectives on Approximation and Sampling Theory. Birkhauser, Cham.
- [4] Aral, A. (2001). On convergence of singular integrals with non-isotropic kernels. Commun. Fac. Sci. Univ. Ank. Ser. A1, 50, 88–98.
- [5] Bardaro, C., Karsli, H. and Vinti, G. (2008). On pointwise convergence of linear integral operators with homogeneous kernels. Integral Transforms Spec. Funct., 19, 429–439.
- [6] Bardaro, C., Karsli, H. and Vinti, G. (2013). On pointwise convergence of Mellin type nonlinear singular integral operators. Comm. Appl. Nonlinear Anal., 20(2), 25–39.
- [7] Bardaro, C., Mantellini, I. (2014). On Mellin convolution operators: a direct approach to the asymptotic formulae. Integral Transf. Spec. Funct., 25(3), 182–195.
- [8] Butzer, P. L., Jansche, S. (1997). A direct approach to the Mellin transform. J. Fourier Anal. Appl., 3(4), 325–376.
- [9] Butzer, P. L., Jansche, S. (1998). The finite Mellin Transform, Mellin-Fourier series and the Mellin-Poisson summation formula. Rend Circ Mat Palermo, 2:52, 55–81.

4th International Conference: Constructive Mathematical Analysis

Mid p -Summable Sequences and Applications to Lipschitz Operators

Ayşegül Keten Çopur

Department of Mathematics and Computer Science, Necmettin Erbakan University, Konya, Türkiye

aketen@erbakan.edu.tr

key-words: Banach space sequences, Lipschitz operators, summing operators.

Abstract:

Mid p -summable sequences were introduced by Karn and Sinha [3]. This notion naturally emerges as an extension of bounded sets within the framework of p -summability. As a new type of summability for vector-valued sequences, mid p -summability has attracted significant attention in the literature and has been studied extensively; see, for example, [2, 1, 5, 4].

Let E be a Banach space and let $1 \leq p < \infty$. A sequence (x_n) in E is said to be *norm p -summable* if $\sum_{n=1}^{\infty} \|x_n\|^p < \infty$, and it is called *weakly p -summable* if $\sum_{n=1}^{\infty} |x^*(x_n)|^p < \infty$ for all $x^* \in E^*$. Every norm p -summable sequence is weakly p -summable; however, the converse does not hold in general and is valid only when E is finite-dimensional. The spaces $\ell_p(E)$ and $\ell_p^w(E)$, consisting of norm and weakly p -summable sequences, respectively, are Banach spaces when endowed with their natural norms. A sequence $(x_j)_{j=1}^{\infty}$ in a Banach space E is called *mid- p -summable* if, for every sequence $(x_n^*)_{n=1}^{\infty} \in \ell_p^w(E^*)$, one has $((x_n^*(x_j))_{j=1}^{\infty})_{n=1}^{\infty} \in \ell_p(\ell_p)$. The collection of all such sequences is denoted by $\ell_p^{\text{mid}}(E)$. This notion provides an intermediate structure between classical summability concepts, as reflected in the inclusions $\ell_p(E) \subseteq \ell_p^{\text{mid}}(E) \subseteq \ell_p^w(E)$.

In this work, a new class of Lipschitz operators is introduced by means of mid p -summable sequences, and several fundamental properties of this class are investigated. Moreover, further results obtained via mid p -summable sequences in the setting of Lipschitz operators are presented.

References

- [1] Botelho, G., Campos, J. R., and Santos, J. (2017). Operator ideals related to absolutely summing and Cohen strongly summing operators. *Pac. J. Math.*, 287(1), 1–17.
- [2] Fourie, J. H., Zeekoei, E. D. (2016). Classes of sequentially limited operators. *Glasg. Math. J.*, 58(3), 573–586.
- [3] Karn, A., Sinha, D. (2014). An operator summability of sequences in Banach spaces. *Glasg. Math. J.*, 56, 427–437.
- [4] Maepa, S. M., Ndumba, B. C. (2024). On mid (p, r) -compact operators. *Quaestiones Mathematicae*, 47(9), 1935–1956.
- [5] Philip, A., Baweja, D. (2023). Ideals of mid p -summing operators: a tensor product approach. *Adv. Oper. Theory*, 8(28).

4th International Conference: Constructive Mathematical Analysis

Numerical Spectral Analysis of a Chafee–Infante Type Equation on Unbounded Domains

Hadil Khenissa¹, Nouria Arar²

¹ Laboratory of Applied Mathematics and Didactics (MAD), École Normale Supérieure de Constantine, Algeria,

² Mathematics and Decision Sciences Laboratory (LAMASD), Université Frères Mentouri Constantine 1, Algeria

khenissa.hadil@ensc.dz, arar.nouria@umc.edu.dz

key-words: Spectral method, Chafee–Infante equation, rational approximation, unbounded domains, weighted Sobolev spaces, Arar–Boulmezaoud functions.

Abstract:

This paper presents a rational spectral method for the numerical approximation of a nonlinear Chafee–Infante type equation posed on the whole real line. The proposed approach relies on the Arar–Boulmezaoud basis functions, which are specifically adapted to unbounded domains and naturally incorporate decay properties at infinity. A weak variational formulation is established in a weighted Sobolev framework, and the resulting nonlinear Galerkin system is discretized using spectral techniques combined with Gauss-type quadrature formulas. The semi-discrete system is integrated in time through a suitable numerical scheme, and the nonlinear term is treated using an iterative procedure. Numerical experiments are performed to validate the accuracy, stability, and spectral convergence of the proposed method on large computational domains.

References

- [1] Arar, N., Boulmezaoud, T. Z. (2013). Eigenfunctions of a weighted Laplace operator in the whole space. *J. Math. Anal. Appl.*, 400, 161–173.
- [2] Boulmezaoud, T. Z., Arar, N., Kerdid, N. and Kourta, A. (2016). Discretization by rational and quasi-rational functions of elliptic problems in the whole space. *ESAIM: M2AN*, 50, 263–288.
- [3] Canuto, C., Hussaini, M. Y., Quarteroni, A. and Zang, T. A. (2006). *Spectral Methods: Fundamentals in Single Domains*. Springer, Berlin.
- [4] Chafee, N., Infante, E. (1974). A bifurcation problem for a nonlinear partial differential equation of parabolic type. *Appl. Anal.*, 4, 17–37.

4th International Conference: Constructive Mathematical Analysis

On Exponential-Type Sampling Series: Upper Bounds of Operator Differences

Sadettin Kursun

¹ Department of Information Security Technology, Izmir Katip Celebi University, Izmir, Türkiye

² Constructive Mathematical Analysis Research Laboratory (CMARL), Selcuk University, Konya, Türkiye

sadettinkursun@yahoo.com

key-words: Exponential-type sampling series, upper bounds for operator differences, Kantorovich-type operators.

Abstract:

In this talk, we present results that provide upper bounds for operator differences in logarithmic weighted spaces of functions. To this end, we consider the following operators: generalized exponential sampling operators [2] and their Kantorovich [3] and generalized Kantorovich forms [1]. Finally, we also discuss several kernel examples that support our theoretical results. It is worth emphasizing that the present study adopts a direct approach within the framework of Mellin analysis for the processes under consideration.

References

- [1] Aral, A., Acar, T. and Kursun, S. (2022). Generalized Kantorovich forms of Exponential Sampling Series. Anal. and Math. Physics, 12(2).
- [2] Bardaro, C., Faina, L. and Mantellini, I. (2017). A Generalization of the Exponential Sampling Series and its Approximation Properties. Math. Slovaca, 67(6), 1481-1496.
- [3] Kumar, S., Bajpeyi, S. (2020). Direct and Inverse Results for Kantorovich Type Exponential Sampling Series. Results Math, 75, 119.

4th International Conference: Constructive Mathematical Analysis

A Mixture Model Using Unit Type Lindley Distributions

Coşkun Kuş¹, Sümeyra Sert¹

¹ Department of Statistics, Selçuk University, Konya, Türkiye

coskun@selcuk.edu.tr, sumeyra.sert@selcuk.edu.tr

key-words: EM algorithm, unit-Lindley distribution, mixture model.

Abstract:

This paper proposes a parsimonious three-parameter mixture model for data supported on the unit interval. The model combines two structurally different unit-Lindley transformations arising from the same Lindley parent distribution. Its novelty lies not in the use of a finite mixture itself, but in bringing these two complementary Lindley-based transformations together within a simple bounded model. This construction preserves the simplicity of one-parameter components while allowing symmetric, asymmetric, sharply peaked, and bimodal shapes. Closed-form expressions are obtained for the density and distribution functions, and several distributional properties are derived, including moments and stress-strength reliability. For the basic three-parameter model, parameter estimation is carried out by the Expectation-Maximization algorithm, and large-sample inference is based on the observed-data information matrix obtained through the missing-information principle. The main empirical application to piped-water access proportions shows that the proposed model can improve likelihood-based fit while yielding interpretable latent profiles. A limited covariate-dependent example is also included only as an illustrative extension.

References

- [1] Louis, T. A. (1982). Finding the observed information matrix when using the EM algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 44(2), 226–233.
- [2] Mazucheli, J., Menezes, A. F. B. and Chakraborty, S. (2019). On the one parameter unit-Lindley distribution and its associated regression model for proportion data. *Journal of Applied Statistics*, 46(4), 700–714.
- [3] Mazucheli, J., Bapat, S. R. and Menezes, A. F. B. (2020). A new one-parameter unit-Lindley distribution. *Chilean Journal of Statistics*, 11(1), 53–67.
- [4] Ng, H. K. T., Chan, P. S. and Balakrishnan, N. (2002). Estimation of parameters from progressively censored data using EM algorithm. *Computational Statistics and Data Analysis*, 39(4), 371–386.

4th International Conference: Constructive Mathematical Analysis

Convergence Domains of Branched Continued Fraction Expansions of Ratio $H_4(a, d+1; c, d; \mathbf{z})/H_4(a+1, d+1; c, d+1; \mathbf{z})$

Ilona-Anna Lutsiv

Department of Mathematical and Functional Analysis, Vasyl Stefanyk Carpathian National University, Ivano-Frankivsk, Ukraine

lutsiv.ilona@gmail.com

key-words: Branched continued fraction, Horn hypergeometric function, convergence.

Abstract:

The Horn's hypergeometric function H_4 is defined by (see [2])

$$H_4(a, b; c, d; \mathbf{z}) = \sum_{k, l=0}^{\infty} \frac{(a)_{2k+l} (b)_l}{(c)_k (d)_l} \frac{z_1^k}{k!} \frac{z_2^l}{l!}, \quad |z_1| < p, \quad |z_2| < q,$$

where a, b, c , and d are complex numbers herewith $c, d \notin \{0, -1, -2, \dots\}$, $(\cdot)_n$ is the Pochhammer symbol, p and q are positive numbers such that $4p = (q-1)^2$ herewith $q \neq 1$, $\mathbf{z} = (z_1, z_2) \in \mathbb{C}^2$.

From Theorem 1 in [1], under the conditions that $b = d+1$ and $(ij)_0 = (1, 2)$, we get that the function

$$\frac{H_4(a, d+1; c, d; \mathbf{z})}{H_4(a+1, d+1; c, d+1; \mathbf{z})} \quad (2)$$

has a formal branched continued fraction expansion of the form

$$1 - \frac{d-a}{d} z_2 - \frac{h_1 z_1}{1-z_2} - \frac{h_2 z_1}{1-z_2} - \frac{h_3 z_1}{1-z_2} - \dots, \quad (3)$$

where

$$h_1 = \frac{2(a+1)}{c}, \quad h_k = \frac{(2c-a+k-3)(a+k)}{(c+k-2)(c+k-1)}, \quad k \geq 2.$$

We consider the problem of establishing the convergence criteria of the branched continued fraction expansion 3 of the ratio 2. To solve it, the technique of expanding the domain of convergence of the branched continued fraction from the known small domain of convergence to a wider domain of convergence is used. Under various conditions on the parameters of the Horn hypergeometric function H_4 , we study different domains of convergence in various unbounded domains of the space $\mathbb{C}^2$.

References

- [1] Antonova, T., Dmytryshyn, R., Lutsiv, I.-A. and Sharyn, S. (2023). On some branched continued fraction expansions for Horn's hypergeometric function $H_4(a, b; c, d; z_1, z_2)$ ratios. Axioms, 12(3), 299.
- [2] Horn, J. (1931). Hypergeometrische Funktionen zweier Veränderlichen. Math. Ann., 105, 381-407.

4th International Conference: Constructive Mathematical Analysis

Pseudospectra in Banach Algebras and Banach Jordan Algebras

Abdelaziz Maouche

Department of Mathematics, Sultan Qaboos University, Oman

maouche@squ.edu.om

key-words: Banach Jordan algebra, quadratic pseudospectrum, numerical range.

Abstract:

This paper provides a study of pseudospectra in Banach algebras and Banach Jordan algebras, systematically transitioning from the classical associative setting to the intrinsically Jordan quadratic framework. Classical pseudospectra are defined via the resolvent, while the Jordan analogue, the quadratic pseudospectrum $\sigma_\epsilon^{(2)}(a)$, is formulated using the quadratic operator U_x which intrinsically governs invertibility in Jordan algebras. Leveraging Moreno's localization theorem which embeds singly generated Jordan subalgebras into full associative commutative Banach subalgebras we establish a universal inclusion $\sigma_\epsilon^{(2)}(a) \subseteq \sigma_\epsilon(a)$ valid for all Banach Jordan algebras. For self-adjoint elements in JB- and JB^* -algebras, we recover exact equivalence,

$$\sigma_\epsilon^{(2)}(a) = \{\lambda : \text{dist}(\lambda, \sigma_A(a)) \leq \epsilon\}$$

and derive direct and inverse quadratic pseudospectral mapping theorems under local injectivity. Moreover, we demonstrate that the numerical range controls quadratic resolvent growth, yielding Kreiss-type estimates and pseudospectral radius bounds in the JB^* -setting. A critical assessment clarifies that exact equality is a C^* -type phenomenon rather than a universal feature, and the numerical range control for general non-self-adjoint elements remains an open problem. Overall, this synthesis validates the quadratic pseudospectrum as the natural stability object in Jordan theory, structurally compatible with classical pseudospectral analysis in all C^* -related contexts.

References

- [1] Kulkarni, S. H., Krishnan, A. (2016). Pseudospectra of elements of reduced Banach algebras. IWOTA.
- [2] Maouche, A. (2026). Pseudospectra in Banach Jordan Algebras. Mathematics, 14(5), 826.
- [3] Martínez Moreno, J. (1991). Holomorphic functional calculus in Jordan-Banach algebras. Ann. Sci. Univ. Blaise Pascal Clermont-Ferrand II, 97, Math. 27, 125-134.

4th International Conference: Constructive Mathematical Analysis

Korovkin and Voronovskaya Results for Positive Bivariate Operators under Generalized Convergence

Francisco-Javier Martínez-Sánchez¹, Pedro Garrancho¹ and Daniel Cárdenas-Morales¹

¹ Departamento de Matemáticas, Universidad de Jaén, Campus Las Lagunillas 23071 Jaén, Spain

fjsanche@ujaen.es, pgarran@ujaen.es, cardenas@ujaen.es

key-words: Korovkin, Voronovskaya, bivariate operator.

Abstract:

This talk is based on a natural continuation of papers [1] and [2]. In [1], the author deals with a qualitative Korovkin-type approximation result about the convergence of sequences of functions defined by means of linear operators, where the limit of the approximation process is the image of a function under certain differential operator, and where a general notion of convergence is utilized. Then, the study is completed by analyzing conditions that guarantee the existence of an asymptotic formula. On the other hand, in [2], the authors prove the corresponding quantitative result, and study the saturation class with the help of that asymptotic formula. While both previous papers were written under a univariate setting, the present one goes into the bivariate case.

The main achievement of this work comes from the fact of considering a general notion of convergence, as it was also the case with [1, 2]. There is a long list of papers where different notions of convergence of double sequences of functions are moved to the setting of approximation theory. Particular double sequences of linear operators are analyzed under particular concepts of convergence, and then, the usual topics within that theory are studied. The framework that we set in this paper, and the results we prove, not only cover all those particular situations, and many previous results in the subject, but provide as well a direct way to deal with further ones that may arise in the future.

References

- [1] Garrancho, P. (2019). A general Korovkin result under generalized convergence. *Constr. Math. Anal.*, 2(2): 81-88.
- [2] Garrancho, P., Martínez-Sánchez, F.J. and Cárdenas-Morales, D. (2024). Approximation rate and saturation under generalized convergence. *Mathematical Foundations of Computing*, 7(1): 148-157.

4th International Conference: Constructive Mathematical Analysis

Bridging Extending and Supplemented Modules: Towards a Unified Framework in Module Theory

Zahra Moaed Abd Al-Majeed¹, Zainab Raad Shaker², Tamara Wathib Mohammed³
and Sura Khaleel Jawad⁴

¹ Department of Chemical Engineering and oil Pollution, College of Chemical Engineering, University of Technology- Iraq, Baghdad, Iraq

² Department of Registration and Students Affairs, University of Technology- Iraq, Baghdad, Iraq

³ Department of Mathematics, College of Science, University of Baghdad, Baghdad, Iraq

⁴ Smart Distinguished Girls Secondary School, Baghdad, Iraq

Zahra.m.abdalmajeed@uotechnology.edu.iq, zainab.r.shaker@uotechnology.edu.iq
tamarah.w@sc.uobaghdad.edu.iq, surakhaleel55@gmail.com

key-words: Extending modules, supplemented modules, bridging modules.

Abstract:

This paper presents a unified theoretical framework connecting extending modules, supplemented modules, and bridging modules. Its main objective is to rigorously investigate their structural relationships, identify underlying similarities, and establish homological conditions under which these classes of modules can be considered equivalent in a categorical sense.

The study adopts a pure mathematical approach based on formal reasoning through lemmas and theorems, supported by explicit constructions and counterexamples. It further highlights new algebraic relations in connection with classical results while addressing existing conceptual gaps.

The proposed framework is illustrated through examples involving $\mathbb{N}$ -modules, Artinian rings, and semisimple rings, demonstrating both its applicability and limitations. A key result shows that, under suitable structural and homological conditions, extending and supplemented modules arise as special cases of bridging modules.

This unification provides deeper insight into module decomposition, exact sequences, and classification theory within homological algebra, offering potential extensions for categorical and computational approaches.

References

- [1] Alahmadi, A., Facchini, A. and Pagliuca, F. (2021). Some natural additive decompositions of elements in bimodules. *Journal of Pure and Applied Algebra*, 226(1), 106806.
- [2] Al-Saadi, S. A., Al-Rubaye, A. A. (2020). -s-extending modules. *Ibn Al-Haitham Journal for Pure and Applied Sciences*, 33(3), 80–92
- [3] Bai, Z., Ma, J. and Wang, Y. (2025). On the annihilator variety of a highest weight module for classical Lie algebras. *Journal of the London Mathematical Society*, 112(2), 1–23.
- [4] Becerril, V. (2025). Gorenstein $\mathcal{FP}_n$ -flat modules and weak global dimensions. *Algebras and Representation Theory*, 28(4), 1073–1091.
- [5] Tattar, A. (2020). Torsion pairs and quasi-abelian categories. *Algebras and Representation Theory*, 24(6), 1557–1581.

4th International Conference: Constructive Mathematical Analysis

A θ -Scheme for Complex Nonlinear Klein-Gordon Equations in the Nonrelativistic Limit: Stability Analysis and Numerical Results

Arezoo Latif Altojar¹, Reza Mokhtari^{1,2}

¹ Department of Mathematical Sciences, Isfahan University of Technology, Isfahan, Iran

² Department for Mathematics and Scientific Computing, University of Graz, Graz, Austria

a.latif@math.iut.ac.ir, mokhtari@iut.ac.ir, reza.mokhtari@uni-graz.at

key-words: θ -scheme, complex nonlinear Klein-Gordon equation, stability analysis.

Abstract:

The nonlinear Klein-Gordon equation (NKGE) stands as a fundamental model in relativistic quantum mechanics for describing spin- $\frac{1}{2}$ particles, and it can be regarded as a relativistic version of the Schrödinger equation [2]. Beyond its theoretical origins, the NKGE has emerged as a pivotal tool in nonlinear optics and plasma physics, where it governs the evolution of robust wave structures and complex multi-scale phenomena in the nonrelativistic limit regime [1]. In this study, we investigate the NKGE within the nonrelativistic limit regime, formulated as

$$\varepsilon^2 \partial_t^2 u = \Delta u - \frac{1}{\varepsilon^2} u - \lambda |u|^2 u, \quad x \in \Omega \subseteq \mathbb{R}, \quad 0 < t \leq T, \quad (1)$$

where u is a complex-valued unknown function and $0 < \varepsilon \ll 1$ denotes a small scaling parameter. Here, $\lambda \neq 0$ is a real constant independent of ε , and T represents a fixed final time [2]. Furthermore, some suitable initial and boundary conditions must be specified for Eq. (1). This regime is characterized by rapid oscillations in both time and space, which present significant challenges for standard numerical approximations. While existing literature often struggles with the computational cost of nonlinear iterations or the rigorous analysis of complex-valued solutions, our work addresses these issues by decoupling the real and imaginary components and employing an efficient linearization technique. We propose a versatile θ -scheme for $\theta \in [0, 1]$, which naturally incorporates the explicit method ($\theta = 0$), the Crank-Nicolson (CN) method ($\theta = 1/2$), and the implicit method ($\theta = 1$) as special cases. A core contribution of this research is a deep theoretical investigation used to derive conditional stability bounds for the entire range of θ , ensuring the reliability of the method across different parameter regimes. A mild condition mentioned in our theorem does not limit the applicability of the method. Numerical experiments, conducted using the discrete L_2 norm, are provided to evaluate the performance of the proposed schemes. As usual, the explicit and implicit schemes provide first-order temporal accuracy. However, the CN scheme exhibits first-order accuracy in time, contrary to the usual second-order behavior. To recover second-order temporal convergence, we propose a three-level θ -scheme.

References

- [1] Bao, W., Dong, X. (2012). Analysis and comparison of numerical methods for the Klein-Gordon equation in the nonrelativistic limit regime. *Numerische Mathematik*, 120(2), 189-229.
- [2] Cui, M., Li, Y. (2025). Energy-conservative finite difference method for the coupled nonlinear Klein-Gordon equation in the nonrelativistic limit regime. *International Journal of Numerical Analysis and Modeling*, 22(2), 246-267.

4th International Conference: Constructive Mathematical Analysis

Convergence and Smoothness of the Mehler-Fock Transform and Application to Overdetermined Boundary Value Problems in $\mathbb{H}^n \times \mathbb{R}$

Filippo Morabito

Università di Udine, Dipartimento Politecnico Ingegneria e Architettura, Via delle Scienze 206, Udine, Italy

filippo.morabito@uniud.it

key-words: Overdetermined problem, integral transform, elliptic pde.

Abstract:

Let $\Omega \subset \mathbb{R}^n$ denote an unbounded smooth domain. Berestycki, Caffarelli, Nirenberg in [1] studied the following boundary value problem

$$\begin{cases} \Delta u + f(u) = 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \\ \frac{\partial u}{\partial \nu} = \text{const} & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where f is a Lipschitz function and ν is the outward unit normal vector field.

They considered domains bounded by the graph of a C^2 Lipschitz function F :

$$\Omega := \{x \in \mathbb{R}^n \mid x_n \geq F(x_1, x_2, \dots, x_{n-1})\}.$$

More precisely, if for any $p \in \mathbb{R}^{n-1}$, uniformly, $\lim_{x \in \mathbb{R}^{n-1}, |x| \rightarrow \infty} (F(x+p) - F(x)) = 0$ and f satisfies some hypotheses, then if there exists a bounded positive solution of (1), the authors showed that Ω is a half-space, that is F is a constant function.

In the same work the authors stated the following conjecture: if Ω is a smooth domain such that $\mathbb{R}^n \setminus \Omega$ is connected and there exists a bounded positive solution of (1) for some Lipschitz function f , then Ω is a half-space, a ball, the complement of a ball, or a circular-cylinder-type domain: $\mathbb{R}^j \times B$ being B a ball. Even though the conjecture is known to be false in the general case, the case when the domain is an epigraph has attracted special attention because of its relation with De Giorgi and Bernstein conjectures.

Overdetermined elliptic problems have been studied in the case when the ambient manifold is a Riemannian manifold as well: the unit sphere, the hyperbolic space, $\mathbb{H}^N \times \mathbb{R}$ and $\mathbb{S}^N \times \mathbb{R}$.

In [2] we showed that under the hypotheses $F \in C^{2,\alpha}$, $f \in C^{1,\alpha}$, $f'(0) \leq 1/4$, there exist unbounded domains Ω (bounded by the vertical graph of a nonconstant function) in $\mathbb{H}^2 \times \mathbb{R}$ for which (1) admits a positive solution. Such domains and the corresponding solutions are obtained by perturbing the half-space $\mathbb{H}^2 \times \mathbb{R}_0^+$ and the solution to $u''(z) + f(u(z)) = 0$ such that $u(0) = 0$, $u(z) \rightarrow \pm 1$ as $z \rightarrow \pm\infty$. The proof is based on the use of the Mehler-Fock transform. In other terms the Berestycki-Caffarelli-Nirenberg conjecture is false in the case when the functions f, F have derivatives which are Hölder functions.

In our recent work we prove the same result in $\mathbb{H}^n \times \mathbb{R}$.

References

- [1] Berestycki, H., Caffarelli, L. A. and Nirenberg, L. (1997). Monotonicity for Elliptic Equations in unbounded Lipschitz domains. Comm. Pure Applied Math., Vol. L, 1089-1111.
- [2] Morabito, F. Overdetermined boundary value problems on unbounded domains of $\mathbb{H}^2 \times \mathbb{R}$. submitted.

4th International Conference: Constructive Mathematical Analysis

A Constructive Definition of the Improper Riemann Integral on a Separable Banach Space

Timothy Myers

Mathematics Department, Howard University, Washington DC, United States

timyers@howard.edu

key-words: Improper Riemann integral, Lebesgue integral, separable Banach space.

Abstract:

The goal of this talk is to construct an improper Riemann integral on a separable Banach space which possesses all of the fundamental properties of the improper Riemann integral on $\mathbb{R}^n$. Let $\mathcal{B}$ represent a separable Banach space. Using an idea from Yamasaki [3], the paper [1] presents a proof that $\mathcal{B}$ has an isomorphic, isometric embedding in $\mathbb{R}^\infty = \mathbb{R} \times \mathbb{R} \times \cdots$. In this work we will use this embedding to define an improper Riemann integral on $\mathcal{B}$ that extends the definition of the Riemann integral on a special rectangle in $\mathcal{B}$ as discussed in [2]; which makes the derivations of most of its properties virtually identical to those of its finite-dimensional analogue.

Similar to the Lebesgue integral on $\mathcal{B}$, this improper Riemann integral is constructive in the sense that it equals a limit of improper Riemann integrals on $\mathbb{R}^n$ as $n \rightarrow \infty$. We will use this convergence to extend some well-known probability density functions to $\mathcal{B}$ and define a Fourier transform on $\mathcal{B}$. This talk will conclude by discussing some ways in which this improper Riemann integral advances the theory of Lebesgue integration on $\mathcal{B}$.

References

- [1] Gill, T.L., Myers, T. (2019). Constructive Analysis on Banach Spaces. *Real Analysis Exchange*, 44, 1-36.
- [2] Myers, T. A. (2025). Constructive Definition of the Riemann Integral on a Separable Banach Space $\mathcal{B}$ and its Relationship to the Lebesgue Integral on $\mathcal{B}$. Part I, 2025 Proceedings of International Conference: Constructive Mathematical Analysis, 131-150.
- [3] Yamasaki, Y. (1980). Translationally Invariant Measure on the Infinite-dimensional Vector Space. *Publications of the Research Institute for Mathematical Sciences*, 16(3), 693-720.

4th International Conference: Constructive Mathematical Analysis

On the Representation of Lauricella-Saran Hypergeometric Functions F_M and Their Ratios in Special Cases by Branched Continued Fractions

Ivan Nyzhnyk

Department of Mathematical and Functional Analysis, Vasyl Stefanyk Carpathian National University, Ivano-Frankivsk, Ukraine

ivan.nyzhnyk.19@pnu.edu.ua

key-words: Hypergeometric function, branched continued fraction, analytic function, approximation by rational functions, convergence, analytic continuation.

Abstract:

The problem of approximating Lauricella-Saran hypergeometric functions F_M and their ratios in special cases by a special family of functions – branched continued fractions – is considered. Under certain conditions of the parameters of the Lauricella-Saran hypergeometric functions F_M , the domains of analytic continuation of the above-mentioned functions are established using their expansions into branched continued fractions, the elements of which are polynomials of three complex variables. Numerical experiments allowed us to evaluate the practical efficient approximating analytic functions by branched continued fractions.

References

- [1] Dmytryshyn, R., Nyzhnyk, I. (2025). On the approximation of Lauricella-Saran's hypergeometric functions F_M and their ratios by branched continued fractions. *Dolomites Res. Notes Approx.*, 18(1), 106–117.
- [2] Lauricella, G. (1893). Sulle funzioni ipergeometriche a più variabili. *Rend. Circ. Matem.*, 7, 111–158.
- [3] Nyzhnyk, I., Dmytryshyn, R. and Antonova, T. (2025). On branched continued fraction expansions of hypergeometric functions F_M and their ratios. *Modern Math. Methods*, 3(1), 1–13.
- [4] Saran, S. (1954). Hypergeometric functions of three variables. *Ganita*, 5, 77–91.

4th International Conference: Constructive Mathematical Analysis

A Randomized Shifted Finite Volume Method to Break Resonance Locking in 1D Homogenization

Bienvenu Ondami

Université Marien Ngouabi, Faculté des Sciences et Techniques, Brazzaville, Congo

bienvenu.ondami@ummng.cg

key-words: Homogenization, finite volume method, resonance locking, random shift, discontinuous coefficients, continuous coefficients.

Abstract:

We investigate the numerical approximation of a one-dimensional elliptic homogenization problem with highly oscillatory periodic coefficients. When the mesh size h is an integer multiple of the period ε ($h/\varepsilon \in \mathbb{N}$), the classical finite volume method suffers from resonance locking: the error does not decrease under mesh refinement. We propose a simple non-intrusive fix: randomly shift the evaluation points of the coefficient at each interface by a small amount uniformly distributed in $[0, h]$. This infinitesimal perturbation breaks the artificial periodicity of the discrete operator. Numerical experiments are performed for both discontinuous (two-phase media with contrasts up to 100) and continuous ($a(y) = 1 + 0.5 \sin 2\pi y$) coefficients, always keeping $h = \varepsilon$. For discontinuous coefficients, the classical pointwise scheme locks (error $\sim 5.5 \times 10^{-2}$), while harmonic averaging (requiring knowledge of discontinuities) achieves second-order accuracy (7.8×10^{-5} at $N = 2560$). Our random shift restores first-order convergence and reduces the error by factors of 30–60 compared to the locked scheme, making it a robust alternative when microstructural details are unknown. For continuous coefficients, harmonic averaging gives no advantage; the random pointwise method again yields first-order convergence and reaches an error of 7.6×10^{-4} at $N = 4096$, which is a gain of 25.5 over the locked classical scheme. The method is trivial to implement, requires no microstructure knowledge, and extends directly to higher dimensions. This randomized shift thus provides a cheap and effective remedy against resonance locking in multiscale computations. The model problem is

$$-\frac{d}{dx} \left(a\left(\frac{x}{\varepsilon}\right) \frac{du}{dx} \right) = 1, \quad x \in (0, 1), \quad u(0) = u(1) = 0, \quad (1)$$

with a periodic. For $h = \varepsilon$ the classical scheme reduces to a constant-coefficient discrete operator, explaining the locking. The random shift introduces variation and restores convergence.

References

- [1] Ondami, B. (2016). Approximation of a second-order elliptic equation with discontinuous and highly oscillating coefficients by finite volume methods. *J. Math. Res.*, 8(6), 34–44.
- [2] Ondami, B. (2015). The effect of numerical integration on the finite element approximation of a second order elliptic equation with highly oscillating coefficients. *J. Interpol. Approx. Sci. Comput.*, (2), 128–136.

4th International Conference: Constructive Mathematical Analysis

Fixed Point Theorems in Digital Metric Spaces and Applications in Fractal Image Compression

Aybala Sevde Özkapu¹, Özlem Acar¹

¹ Department of Mathematics, Faculty of Science, Constructive Mathematical Analysis Research Laboratory (CMARL), Selçuk University, Konya, Türkiye

aybalasevdeozkapu@yahoo.com, acarozlem@ymail.com

key-words: Digital metric space, fractal image compression, vector-valued metric.

Abstract:

Fixed point theory has long been recognized as a powerful tool across various fields, from functional analysis to computer science [1, 3]. However, when problems are considered in a digital setting, such as image analysis, the continuous framework must be adapted to discrete structures and adjacency relations [1]. Digital metric spaces provide a natural environment for extending fixed point results to model such discrete data [1]. This seminar will comprehensively explore next-generation contraction mappings developed within digital metric spaces, specifically focusing on digital F -contractions [1], rational type common fixed point theorems [3, 4], and θ -contractions [2]. Moving beyond traditional scalar approaches, the concept of vector-valued digital metric spaces will be introduced to address multi-objective optimization problems, such as simultaneously measuring pixel intensity and structural gradients [4, 2]. In addition to the theoretical foundations, the practical efficacy of these mathematical frameworks will be demonstrated through their applications in fractal image compression [3, 4, 2]. Specifically, the seminar will showcase how fractal-based models around the digital Sierpinski triangle [3] and the novel Adaptive Digital F-Fractal Compression (ADFFC) algorithm [4] successfully optimize the trade-off between image fidelity and compression ratio using fixed point iterations.

Acknowledgements: The first author has been supported by Scientific and Technological Research Council of Turkey (TUBITAK)/ 2224-B-Grant Program for Participation in Scientific Meetings within the Country. The author thanks to TUBITAK for their supports.

References

- [1] Acar, Ö., Han S.E., and Özkapu, A. S. (2026). New Results on a Digital Metric Space via an F -Contraction. Submitted.
- [2] Acar, Ö., Özkapu, A. S. (2026). θ -Contractions on Vector-Valued Digital Metric Spaces and Image Compression. Submitted.
- [3] Özkapu, A. S., Acar Ö. (2026). New common fixed point results in digital metric space. *Demonstratio Mathematica*, 59(1): 20250208, 1-10.
- [4] Özkapu, A. S. (2026). Generalized Digital F -Contractions and Adaptive Image Compression. Submitted.

4th International Conference: Constructive Mathematical Analysis

Approximation by a Novel Modification of a Mellin-Type Kernel

Firat Ozsarac

Department of Mathematics, Kırıkkale University, Kırıkkale, Türkiye

firat_ozsarac@hotmail.com

key-words: Logarithmic moments, Mellin derivative, Euler-Gamma function.

Abstract:

In this presentation, we express the approximation features for a novel modification of the Mellin-type kernel. Moreover, a Voronovskaya-type theorem is given using the Euler Gamma function and Mellin derivatives.

References

- [1] Bardaro, C., Mantellini, I. (2007). Voronovskaya-type estimates for Mellin convolution operators. *Res. Math.*, 50, 1–16.
- [2] Bardaro, C., Mantellini, I. (2010). A quantitative Voronovskaya formula for Mellin convolution operators. *Mediterr. J. Math.*, 7, 483–501.
- [3] Bardaro, C., Mantellini, I. (2011). A note on the Voronovskaja theorem for Mellin-Fejer convolution operators. *Appl. Math. Lett.*, 24, 2064–2067.
- [4] Bardaro, C., Mantellini, I. (2011). Asymptotic behaviour of Mellin-Fejer convolution operators. *East J. Approx.*, 17, 181–201.
- [5] Butzer, P. L., Jansche, S. (1997). A direct approach to the Mellin transform. *J. Fourier Anal. Appl.*, 3, 325–375.
- [6] Mamedov, R. G. (1991). *The Mellin Transform and Approximation Theory*. Elm, Baku, (in Russian).

4th International Conference: Constructive Mathematical Analysis

Hyperbolic Wavelets, Basis Problem

Margit Pap

Faculty of Sciences, University of Pécs, Pécs, Hungary

papm@gamma.ttk.pte.hu

key-words: Hyperbolic wavelets, basis problem in the disc algebra.

Abstract:

We study the explicit hyperbolic Malmquist–Takenaka system associated with the logarithmic hyperbolic grid in the unit disc. We proved that the corresponding multiresolution levels form a complete model set in the disc algebra $A(\mathbb{D})$; equivalently, the union of the finite resolution spaces is dense in $A(\mathbb{D})$. We then investigate the convergence of the orthogonal projections in the supremum norm. A direct kernel-based approach leads to a multiscale shell decomposition. This yields uniform control of the low-frequency and bounded-width critical-frequency contributions, but also shows that absolute L^1 -kernel estimates are too crude to capture the cancellation present in the high-frequency regime. Motivated by this obstruction, we develop two alternative reductions of the max-norm convergence problem. The first is operator-theoretic: using the interpolation property of the projections and the exact Christoffel–Darboux formula for the detail kernels, we reduce the problem to an off-scale decay estimate for the detail operators $D_j = P_j - P_{j-1}$. The second is sampling-based: using the geometry of the hyperbolic lattice, we reduce convergence to a uniform norming-set inequality for the finite resolution spaces. As a rigorous partial result, we prove max-norm convergence for all functions whose detail contributions are summable in the supremum norm.

References

- [1] Pap, M. (2011). Hyperbolic wavelets and multiresolution in $H^2(\mathbb{T})$. *J. Fourier Anal. Appl.*, 17, 755–776.
- [2] Pap, M. (2015). A Special Voice Transform, Analytic Wavelets, and Zernike Functions. In: P. W. Hawkes (Ed.), *Advances in Imaging and Electron Physics*, New York: Elsevier, Vol. 188, 79–134.
- [3] Pap, M., Schipp, F. (2008). The voice transform on the Blaschke group I. *Pure Math. Appl.*, 17(3-4), 387–395.

4th International Conference: Constructive Mathematical Analysis

A Smooth Lagrangian-Based Neural Network for Solving Semidefinite Programming

Pakkapon Preechasilp¹, Rattanaorn Wangkeeree², Rabian Wangkeeree², Kasamsuk Ungchittrakool², Yirga Abebe Belay^{2,3} and Panatda Boonman⁴

¹ Program in Mathematics, Faculty of Education, Pibulsongkram Rajabhat University, Phitsanulok, Thailand

² Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, Thailand

³ Department of Mathematics, Aksum University, Aksum, Ethiopia

⁴ Faculty of Science and Technology, Rajamangala University of Technology Suvarnabhumi, Phra Nakhon Si Ayutthaya, Thailand

pakkapon.p@psru.ac.th, rattanapornw@nu.ac.th

key-words: Semidefinite programming, neural network, augmented lagrangian smoothing.

Abstract:

In this talk, we introduce a smooth gradient-based neural network for semidefinite programming (SDP) using the augmented Lagrangian smoothing (ALS) mapping. The proposed model is formulated through a continuous energy function whose equilibrium points correspond to the Karush–Kuhn–Tucker (KKT) conditions of the SDP problem. We present the main theoretical results, including well-posedness, Lyapunov stability, global convergence, and local exponential convergence. Numerical experiments on benchmark problems show that the method converges reliably and performs competitively compared with existing smoothing-based neural network approaches. In particular, while the proposed method has comparable accuracy on eigenvalue optimization problems, it gives substantially better results for spectral norm minimization. The framework also introduces a penalty parameter ρ that provides additional flexibility in balancing primal and dual variables.

References

- [1] Nikseresht, A., Nazemi, A. (2022). A smoothing gradient-based neural network strategy for solving semidefinite programming problems. *Network: Computation in Neural Systems*, 33(3–4), 187–213.
- [2] Zhang, R.-J., Liu, X.W. and Dai, Y.H. (2024). IPRSDP: A primal-dual interior-point relaxation algorithm for semidefinite programming. *Computational Optimization and Applications*, 88, 1–36.

4th International Conference: Constructive Mathematical Analysis

Rational Approximation and Meromorphic Continuation

Vasiliy Prokhorov

Department of Mathematics and Statistics, University of South Alabama, Mobile, Alabama, USA

prokhorov@southalabama.edu

key-words: Rational approximation, meromorphic continuation, Walsh table.

Abstract:

New asymptotic results for products of best rational approximation errors taken from several rows of the Walsh table will be presented. These results connect the rate of rational approximation with radii of meromorphic continuation, extending classical theorems of Hadamard and Saff-Gonchar. The proof uses Hankel operators and an AAK-type theorem.

References

- [1] Gonchar, A.A. (1974). On a theorem of Saff. *Mat. Sb. (N.S.)* 94 (136), 1, 152–157.
- [2] Hadamard, J. (1892). Essai sur l'étude des fonctions données par leur développement de Taylor, J. *Math. Pures Appl. (4)* 8, 101–186.
- [3] Prokhorov, V.A. (2026). Hadamard-type asymptotics for products of best rational approximation errors. *Math arXiv:2604.03854*, 1–32.
- [4] Saff, E.B. (1971). Regions of meromorphy determined by the degree of best rational approximation. *Proc. Amer. Math. Soc.* 29, 30–38.

4th International Conference: Constructive Mathematical Analysis

Degenerate Appell Sequences via Caputo-Type Fractional Difference Operator

Stiven Díaz¹, William Ramírez¹, Yamilet Quintana² and Clemente Cesarano³

¹ Department of Natural and Exact Sciences, Universidad de la Costa, Barranquilla, Colombia

² Department of Mathematics, Universidad Carlos III de Madrid, Madrid, Spain

³ Section of Mathematics, International Telematic University Uninettuno, Rome, Italy

`sdiaz47@cuc.edu.co`, `wramirez4@cuc.edu.co`, `yaquinta@math.uc3m.es`,
`c.cesarano@uninettunouniversity.net`

key-words: Caputo operator, Mittag-Leffler function, ∇_λ -Appell sequences.

Abstract:

We introduce the notion of ∇_λ^α -Appell sequences of functions based on the discrete fractional Caputo operator and the degenerate Mittag-Leffler function. We analyze several properties of these function sequences, including explicit expressions for each term and their associated generating functions. In particular, we establish some basic novel properties for the case $\alpha = 1$, in which the corresponding sequences become ∇_λ -Appell polynomial sequences.

It is worth noting that the use of fractional operators and expansions in fractional powers which mimic the structural form of Appell (or Scheffer) sequences is a recent development (cf. [1, 2] and the references therein). To the best of our knowledge, the novelty of this work has been to study an analogous situation incorporating the notion of the degeneracy (cf. [3, 4, 5]).

References

- [1] Caratelli, D., Natalini, P. and Ricci, P. E. (2023). Examples of expansions in fractional powers, and applications. *Symmetry*, 15(9) (2023).
- [2] Caratelli, D., Natalini, P. and Ricci, P. E. (2023). Fractional Bernoulli and Euler numbers and related fractional polynomials—A symmetry in number theory. *Symmetry*, 15(10).
- [3] Cesarano, C., Quintana, Y. and Ramírez, W. (2024). Degenerate versions of hypergeometric Bernoulli–Euler polynomials. *Lobachevskii J. Math.*, 45(8), 3509–3521.
- [4] Costabile, F. A., Longo, E. (2013). Δ_h -Appell sequences and related interpolation problem. *Numer. Algorithm*, 63, 165–186.
- [5] Díaz, S., Ramírez, W., Quintana, Y. and Cesarano, C. (2026). Degenerate Appell sequences via Caputo-type fractional difference operator. To appear in *Rend. Circ. Rend. Circ. Mat. Palermo (II)*.

4th International Conference: Constructive Mathematical Analysis

Inequalities for Complex Polynomials

Poonam Babani Rane¹, Prasanna Kumar¹

¹ Department of Mathematics, BITS Pilani, K K Birla Goa Campus, Goa, India

p20230424@goa.bits-pilani.ac.in, prasannak@goa.bits-pilani.ac.in

key-words: Polynomials, zeros, inequalities.

Abstract:

A result concerning the inequalities for complex polynomials, credited to Rivlin [1], states that for a polynomial $P(z)$ of degree n that has no zeros in the disc $|z| < 1$, the following inequality holds:

$$\max_{|z|=r} |P(z)| \geq \left(\frac{1+r}{2} \right)^n \max_{|z|=1} |P(z)|,$$

for $0 \leq r \leq 1$. Rivlin's inequality has been generalized to encompass discs of radius greater than or equal to 1 by several mathematicians. In this presentation, we discuss more refined and generalized inequalities, of which a specific case would enhance the classical Rivlin's inequality.

References

- [1] Rivlin, T. J. (1960). On the maximum modulus of polynomials. Amer. Math. Monthly, 67(3), 251-253.

4th International Conference: Constructive Mathematical Analysis

Spherical Number Algebras: An Invariant-First Framework from Conceptual Architecture to Symbolic and Numeric Presentations

Mohammad Reza Alaei Jordehi

Independent Researcher, Iran

mr.alaeij@gmail.com

key-words: Spherical number algebras, point-free mathematics, invariant structures.

Abstract:

This paper introduces Spherical Number Algebras (SNA), a new structural framework in pure mathematics founded on the principles of admissibility-first, invariant-first, and point-free reasoning. Rather than assuming primitive objects, fields, numbers, or spatiotemporal substrates, SNA takes admissible assembly-events as its fundamental entities and treats structural admissibility as logically prior to composition. Each admissible event carries an invariant package from which observables, symbolic descriptions, and numerical representations arise as derived projections.

The framework is organized around four guiding principles—admissibility before composition, invariants before observation, articulation before objecthood, and point-free continuity before numeric representation—providing a pre-numeric mathematical foundation in which algebraic and symbolic structures emerge through successive translation layers.

The development proceeds in three parts. Part I establishes the conceptual foundations of admissible assembly-events, invariant packages, and observation as a derived process. Part II develops a point-free pre-numeric formal core based on word rewriting, proving that observable assemblies form canonical quotient structures with well-defined composition, induced invariant projections, and associativity up to observation. Part III introduces symbolic, discrete, and numeric translation layers through coding maps from stabilized admissible events or their observable quotients, where rewriting systems operate on symbolic codes and numerical values arise only as tertiary projections.

Throughout the framework, invariant packages remain primary, while symbolic and numerical descriptions are treated as representational layers rather than ontological primitives. SNA therefore provides an invariant-centered mathematical foundation preceding conventional numeric, metric, probabilistic, or energetic formulations.

References

- [1] Awodey, S. (2010). *Category Theory*, 2nd ed.. Oxford University Press.
- [2] Baader, F., Nipkow, T. (1998). *Term Rewriting and All That*. Cambridge University Press.
- [3] Birkhoff, G. (1967). *Lattice Theory*. 3rd ed., American Mathematical Society.
- [4] Burris, S., Sankappanavar, H. P. (1981). *A Course in Universal Algebra*. Springer.
- [5] Cohn, P. M. (1981). *Universal Algebra*. D. Reidel Publishing Company.
- [6] Davey, B. A., Priestley, H. A. (2002). *Introduction to Lattices and Order*, 2nd ed.. Cambridge University Press.
- [7] Johnstone, P. T. (1982). *Stone Spaces*. Cambridge University Press.
- [8] Mac Lane, S. (1998). *Categories for the Working Mathematician* 2nd ed.. Springer.

4th International Conference: Constructive Mathematical Analysis

Sums as Sets

Anthony J. Ripa

Computer Science, Stony Brook University, Stony Brook, USA

aripa@cs.stonybrook.edu

key-words: Set-theoretic calculus, foundations of calculus, dimensional analysis.

Abstract:

This paper develops a set-theoretic account of calculus in which sums are treated as sets before they are treated as numbers. The motivating contrast is between numerical difference and set difference: while an expression such as $f(x) - f(x - h)$ becomes problematic when the width collapses, the corresponding set-theoretic difference remains meaningful, as in $(a, b] \setminus (a, b) = \{b\}$. This calculus preserves the exact set-theoretic difference instead of first projecting to numbers.

Setup. Let f be a set of tuples over R (for concreteness, $R = \mathbb{R}$ suffices), with last coordinate designated as abscissa. Write $f_{\leq b}$, $f_{< b}$, and $f_{=b}$ for the parts whose abscissa is respectively at most, less than, or equal to b . The basic decomposition is $f_{\leq b} = f_{< b} \sqcup f_{=b}$, hence $f_{\leq b} \setminus f_{< b} = f_{=b}$.

Evaluation. Define evaluation as abscission of f at b as $|^b f := f_{\leq b}$. Further, $|^b_a f := |^b f \setminus |^a f$.

Example. Our running example is simplices: $t_n := \{x \in [0, 1]^n \mid x_1 \leq \dots \leq x_n\}$, and $t := t_1$.

Differential. We define differential as set difference $\Delta^b f := f_{\leq b} \setminus f_{< b}$. Thus, $\Delta^b f = f_{=b}$. In one dimension, a differential is a singleton; in two dimensions, it is a strip, i.e. a one-dimensional part crossed with a singleton. For example, $\Delta^b t = \{b\}$, $\Delta^b t_2 = [0, b] \times \{b\}$, and $\Delta^1 t_n = t_{n-1} \times \{1\}$.

Sums. We define the sum Σf as $(\Sigma f)_{=b} := |^b f \times \Delta^b t$. Equivalently, $\Sigma f = \bigsqcup_b (|^b f \times \{b\})$. We also define $\Sigma^b_a f := |^b_a(\Sigma f)$, and $\Sigma^b f := |^b(\Sigma f)$. For example, $\Sigma^1 t_n = t_{n+1}$. So, linear f makes Σf a triangle ($\frac{\text{square}}{2}$), as expected. Similarly, $\Sigma \Sigma f$ is a triangular pyramid ($\frac{\text{cube}}{6}$), as expected.

Quotient. Cartesian quotient is inverse to Cartesian product: $C = A \times B \implies C/B = A$.

Derivative. We define the derivative to be the Cartesian quotient of an exact differential. $D^b f := \Delta^b f / \Delta^b t$. For example, $D^1 t_n = t_{n-1}$. Further, we define $DF := f \iff \forall b (D^b F = |^b f)$.

Fundamental Theorems of Calculus (FTC).

Part 1: $D\Sigma f = f$. Any anti-sum (f) is the derivative: $F = \Sigma f \implies f = DF$.

Part 2: $\Sigma DF = F$. Any anti-derivative (F) is the sum: $f = DF \implies F = \Sigma f$.

Combining 1 & 2, is the unified fundamental theorem: $f = DF \iff F = \Sigma f$.

Example. For a finite example, let $R = \{1, 2, 3\}$. Let $f = \{(1), (2), (3)\} \subseteq R^1$.

The evaluations are $|^1 f = \{(1)\}$, $|^2 f = \{(1), (2)\}$, and $|^3 f = \{(1), (2), (3)\}$.

The differentials are $\Delta^1 f = \{(1)\}$, $\Delta^2 f = \{(2)\}$, and $\Delta^3 f = \{(3)\}$.

The sum is $\Sigma f = \bigsqcup_b (|^b f \times \{b\}) = \{(1, 1), (1, 2), (2, 2), (1, 3), (2, 3), (3, 3)\}$.

FTC1: $D\Sigma f = f$. E.g., here $\Delta^2(\Sigma f) = \{(1, 2), (2, 2)\}$. $D^2(\Sigma f) = \{(1), (2)\} = |^2 f$.

FTC2: $DF = f \implies$ layer by layer $F = \{(1, 1), (1, 2), (2, 2), (1, 3), (2, 3), (3, 3)\}$, which is Σf .

Summary. We replace infinitesimal or limiting differentials with set-theoretic differentials. A differential is a literal subset; differentiation is Cartesian quotient; and integration is union. Numerical measures are optional representations of structure already present at the level of sets.

References

- [1] Viète, F. (1591). *In Artem Analyticem Isagoge*, Tours: Jamet Mettayer.
https://archive.org/details/bub_gb_BWTyywN39KEC

4th International Conference: Constructive Mathematical Analysis

On the Oscillatory Behavior of Cesàro Means for Fourier-Jacobi Series

Samra Sadiković Hajrić¹, Jasmina Muminović Huremović¹

¹ Department of Mathematics, University of Tuzla, Tuzla, Bosnia and Herzegovina

samra.sadikovic@untz.ba, jasmina.muminovic@untz.ba

key-words: Fourier-Jacobi series, Cesàro means, dyadic increments.

Abstract:

We investigate oscillatory properties of Cesàro means $\{\sigma_n^\delta(f, x)\}_{n \geq 0}$, ($\delta > -1$), for Fourier-Jacobi expansions and their connection with r -variation norms, motivated by recent developments in variation estimates for Fourier series and Carleson-type operators [3]. Using Abel-type transformations and kernel estimates, we derive increment formulas and introduce a near-far decomposition for dyadic increments. We prove that the far contribution is controlled by the maximal partial sum operator, yielding L_p -bounds for dyadic consecutive increment functionals, in the classical mean convergence range for Fourier-Jacobi expansions [2]. We further analyze the kernel structure of the associated dyadic increment operators and identify the near-diagonal region as the principal source of difficulty in obtaining full variation estimates. The results provide a structural framework for future study of variation-norm estimates for Cesàro means of Fourier-Jacobi series.

References

- [1] Chanillo, A., Muckenhoupt, B. (1993). Weak Type Estimates for Cesaro Sums of Jacobi Polynomial Series, Mem. Amer. Math. Soc., 102 , (487), viii–90.
- [2] Guadalupe, J. J. Pérez, M. F. Ruiz, J. and Varona J. L. (1992). Two notes on convergence and divergence a.e. of Fourier series with respect to some orthogonal systems. Proc. Amer. Math. Soc., 116(2), 457-464.
- [3] Oberlin, R. Seeger, A. Tao, T. Thiele, C. and Wright, J. (2012). A Variation Norm Carleson Theorem. J. Eur. Math. Soc., 14(2), 421-464.

4th International Conference: Constructive Mathematical Analysis

Modeling Bounded Data for the Unit Transformed Inverse Weibull Distribution: Estimation and Application

Sule Saglam¹, Kadir Karakaya¹

¹ Department of Statistics, Faculty of Science, Selcuk University, Konya, Türkiye

sulesaglam75@gmail.com, kkarakaya@selcuk.edu.tr

key-words: Bonferroni and Lorenz curves, bounded distribution, quantile function, maximum likelihood estimation, Monte Carlo simulation.

Abstract:

In this study, we propose a new bounded distribution, called as unit transformed inverse Weibull distribution. The proposed model is defined on the unit interval $(0, 1)$ and is derived from the transformed inverse Weibull distribution defined on $(0, \infty)$. The proposed model extends the flexibility of classical bounded distributions and is particularly suitable for modeling proportion data. Several fundamental statistical properties of the unit transformed inverse Weibull distribution are derived, including the probability density, cumulative function and hazard rate distribution functions, quantile function, order statistics, and entropy, etc. The proposed distribution is characterized are further examined through, the Bonferroni and Lorenz curves. Several estimation methods are employed to estimate the unknown model parameters. The performance of these estimators is evaluated through an extensive Monte Carlo simulation study based on bias, mean squared error, mean absolute bias, and mean relative error criteria. An application to a real data set is presented to illustrate the practical relevance and modeling capability of the proposed distribution.

References

- [1] Gong, Q., Zhang, Z., Zeng, L., and Ren, H. (2025). A novel extended inverse Weibull distribution: Statistical analysis and application. Plos one, 20(10).

Reliability Evaluation for the Linear or Circular Consecutive k-out-of-n: G System Using Continuous Time Bayesian Network

Ranganathan Sakthivel¹, Gundala Vijayalakshmi² and Sahadevan Sakthipriya³

¹ Department of Career Development Centre, College of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu, India

² Department of Mathematics, College of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu, India

³ Department of Computing Technology, College of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu, India

sakthivr@srmist.edu.in, vijayalg@srmist.edu.in, sakthips4@srmist.edu.in

key-words: Reliability analysis, continuous time Bayesian Network, consecutive k-out-of-n: G system.

Abstract:

This paper presents an analysis of linear and circular consecutive k-out-of-n:G systems using the continuous-time Bayesian network (CTBN) approach. The main contributions include the development of CTBN models for these systems and the subsequent reliability analysis. Additionally, the paper derives exact expressions for system reliability and mean time to failure (MTTF). Numerical illustrations are presented. It is concluded that the reliability of circular system is more than linear system.

References

- [1] Gökdere, G., Tony Ng, H. K. (2022). Time-dependent reliability analysis for repairable consecutive-k-out-of-n: F system. *Statistical Theory and Related Fields*, 6(2), 139-147.
- [2] Kong, D., Lin, Z., Li, W. and He, W. (2024). Development of an improved Bayesian network method for maritime accident safety assessment based on multiscale scenario analysis theory. *Reliability Engineering and System Safety*, 251, 110344.
- [3] Kuo, W., Zhang, W. and Zuo, M. (1990). A consecutive-k-out-of-n: G system: The mirror image of a consecutive-k-out-of-n: F system. *IEEE Transactions on Reliability*, 39(2), 244-253.
- [4] Meng, H., Hu, M., Kong, Z., Niu, Y., Liang, J., Nie, Z. and Xing, J. (2024). Risk analysis of lithium-ion battery accidents based on physics-informed data-driven Bayesian networks. *Reliability Engineering and System Safety*, 251, 110294.
- [5] Sharma, K. K., Krishna, H. (1994). Bayesian reliability analysis of a k-out-of-m system and the estimation of sample size and censoring time. *Reliability Engineering and System Safety*, 44(1), 11-15.
- [6] Tao, H., Jia, P., Wang, X. and Wang, L. (2024). Reliability analysis of subsea control module based on dynamic Bayesian network and digital twin. *Reliability Engineering and System Safety*, 248, 110153.
- [7] Tong, Y. L. (1985). A rearrangement inequality for the longest run, with an application to network reliability. *Journal of Applied Probability*, 22(2), 386-393.
- [8] Xiahou, T., Zheng, Y. X., Liu, Y. and Chen, H. (2023). Reliability modeling of modular k-out-of-n systems with functional dependency: A case study of radar transmitter systems. *Reliability Engineering and System Safety*, 233, 109120.

4th International Conference: Constructive Mathematical Analysis

Modern Iterative Optimization with Monotone Operators and Nonexpansive Mappings

Berat Sert¹, Özlem Acar¹

¹ Department of Mathematics, Faculty of Science, Selçuk University, Konya, Türkiye

beratsert10@hotmail.com, acarozlem@ymail.com

key-words: Halpern iteration, proximal gradient method, strong convergence.

Abstract:

In recent years, convex optimization and fixed point theory have emerged as indispensable tools for solving complex inverse problems in engineering and computer sciences. While the standard proximal gradient method is highly efficient for real-world tasks such as image deblurring and denoising, its application to high-dimensional problems is often hindered by slow convergence rates. To overcome this limitation, inertial extrapolation techniques are widely employed. However, a fundamental theoretical challenge arises in infinite-dimensional Hilbert spaces, where most existing accelerated algorithms rely heavily on the Mann-type iteration, guaranteeing only weak convergence.

In this study, we introduce a novel Halpern-type inertial proximal gradient algorithm that simultaneously achieves acceleration and strong convergence guarantees. We propose a new three-step hybrid iterative scheme that unifies the geometric properties of Halpern anchoring, inertial momentum, and proximal splitting operators. This structural approach elegantly resolves the theoretical paradox between the "shoot forward" nature of inertial extrapolation and the "pull backward" stabilization of Halpern anchoring.

Under very relaxed control conditions on the parameter sequences, we prove the strong convergence of the proposed scheme to an optimal solution of the minimization problem without assuming monotonicity. Furthermore, we extend our generalized optimization theorem to solve the prominent Split Feasibility Problem (SFP) in set intersections. Computational experiments via PSNR and SSIM metrics confirm that our novel algorithm significantly outperforms classical and weakly-convergent inertial algorithms in terms of both operational speed and structural edge preservation in digital image processing.

References

- [1] Combettes, P. L. (1993). The Foundations of Set Theoretic Estimation. IEEE Transactions on Signal Processing, 41(2), 182-208.
- [2] Halpern, B. (1967). Fixed Points of Nonexpanding Maps. Bulletin of the American Mathematical Society, 73(6), 957-961.
- [3] Rockafellar, R. T. (1976). Monotone Operators and the Proximal Point Algorithm. SIAM Journal on Control and Optimization, 14(5), 877-898.

4th International Conference: Constructive Mathematical Analysis

A New Nonparametric k -Sample Test Based on Interval Counts of Order Statistics

Sümeysra Sert¹, Coşkun Kuş¹

¹ Department of Statistics, Selçuk University, Konya, Türkiye

sumeyra.sert@selcuk.edu.tr, coskun@selcuk.edu.tr

key-words: K-sample problem, nonparametric test, order statistics.

Abstract:

In this study, we propose a new nonparametric test statistic for the k -sample problem, which tests the null hypothesis that k independent samples are drawn from the same continuous distribution. The proposed test is an extension of the two-sample precedence-type tests and based on the placement of order statistics. Specifically, the test statistic is constructed by summing the counts of observations from one sample falling into intervals defined by the order statistics of another sample, considered symmetrically across all pairs of groups. Due to the complex dependency structure of the statistic, we employ a permutation-based approach to determine critical values. Monte Carlo simulations are conducted to evaluate the power of the proposed test and compare its performance against the traditional Kruskal-Wallis test under various location and scale shift scenarios. A numerical example is also provided to illustrate the applicability of the proposed test.

References

- [1] Balakrishnan, N., Ng, H. T. (2006). Precedence-type tests and applications. John Wiley and Sons, Vol. 472.
- [2] Kruskal, W.H., Wallis W.A. (1952). Use of ranks in one-criterion variance analysis. J. Am. Stat. Assoc., 260(47), 583-621.

4th International Conference: Constructive Mathematical Analysis

Fixed Point Theorems for Nonlinear Contractive Mappings in Cone-Valued θ -Type Multiplicative Metric Spaces

Pravin Singh¹, Virath Singh¹, Shivani Singh² and Sani Salisu¹

¹ Mathematics, University of Kwazulu-Natal, Durban, South Africa

² Department of Decision Sciences, University of South Africa, Pretoria, South Africa

singhp@ukzn.ac.za, singhv@ukzn.ac.za, singhs2@unisa.ac.za, salisus@ukzn.ac.za

key-words: Banach algebra, contractions, metric.

Abstract:

Fixed point theory plays a fundamental role in nonlinear analysis and has significant applications in differential equations, integral equations, optimization, and applied mathematics. Inspired by recent developments in generalized metric structures, this paper introduces and studies Geraghty-type contractive conditions in the setting of cone θ -type multiplicative metric spaces defined on ordered Banach algebras. By extending the classical notions of θ -metric and multiplicative metric spaces, we construct a framework based on solid multiplicative cones and establish their essential topological and convergence properties. Within this generalized structure, we formulate new fixed point theorems for self-mappings satisfying cone-valued θ -type multiplicative Geraghty contractions, which weaken the traditional Lipschitz condition while preserving the existence and uniqueness of fixed points in complete spaces. The proposed results significantly extend and unify several well-known contraction principles, including those of Banach, Kannan, Chatterjea, and standard Geraghty contractions, under a broader cone-valued multiplicative setting. Furthermore, we develop extensions to higher-dimensional frameworks to enhance applicability in complex nonlinear systems. Illustrative examples are presented to substantiate the theoretical findings and to demonstrate the effectiveness of the introduced approach in generalized analytical environments.

References

- [1] Altomare, F. (2019). Iterates of Markov Operators and Constructive Approximation of Semigroups. *Constr. Math. Anal.*, 2(1), 22-39.
- [2] Aral, A. (2023). Weighted approximation: Korovkin and quantitative type theorems. *Modern Math. Methods*, 1(1), 1-21.
- [3] Karsh, H. (2018). Approximation Results for Urysohn Type Two Dimensional Nonlinear Bernstein Operators. *Constr. Math. Anal.*, 1(1), 45-57.

4th International Conference: Constructive Mathematical Analysis

Dynamic Aspects of Composition Operators on Weighted Holomorphic Function Spaces

Hayri Topal

Department of Mathematics, Van Yuzuncu Yıl University, Van, Türkiye

hayritopal@yyu.edu.tr

key-words: Composition operators, mean ergodic, holomorphic functions.

Abstract:

Let X be a complex Banach space, and let B_X denote the open unit ball. We denote by $H(B_X)$ the space of all holomorphic function $f : B_X \rightarrow \mathbb{C}$. Any continuous and bounded mapping $v : B_X \rightarrow (0, \infty)$ is called a weight. For the weight v defined in this way, we define the following weighted Banach spaces of holomorphic functions on B_X

$$H_v(B_X) = \{f \in H(B_X) : \|f\|_v = \sup_{x \in B_X} v(x)|f(x)| < \infty\}.$$

We investigate the dynamical properties of composition operators acting on the weighted space of holomorphic functions $H_v(B_X)$. By using the asymptotic behavior of the inducing self-maps $\varphi : B_X \rightarrow B_X$, we obtain explicit sufficient criteria ensuring that the associated composition operator is power bounded or uniformly mean ergodic. Several examples are provided to illustrate the applicability of the obtained results. Many researchers have investigated the dynamic properties of composition operators on various spaces of holomorphic functions, and extensive studies on this topic can be found in the literature [1, 2, 3, 4].

References

- [1] Beltrán-Meneu, M. J., Gómez-Collado, M. C., Jordá, E. and Jornet, D. (2016). Mean ergodic composition operators on Banach spaces of holomorphic functions. *J. Funct. Anal.*, 270, 4369–4385.
- [2] García, D., Maestre, M. and Sevilla-Peris, P. (2004). Composition operators between weighted spaces of holomorphic functions on Banach spaces. *Ann. Acad. Sci. Fenn. Math.*, 29, 81–98.
- [3] Jornet, D., Santacreu, D. and Sevilla-Peris, P. (2021). Mean ergodic composition operators on spaces of holomorphic functions on a Banach space. *J. Math. Anal. Appl.*, 500, 125-139.
- [4] Manhas, J. S. (2013). Weighted Composition Operators and Dynamical Systems on Weighted Spaces of Holomorphic Functions on Banach Spaces. *Ann. Funct. Anal.*, 4, 58–71.

4th International Conference: Constructive Mathematical Analysis

Simultaneous Approximation by Bivariate Kantorovich-Type Operators in Weighted Spaces

Metin Turgay

¹ Department of Mathematics, Faculty of Science, Selçuk University, Selçuklu, Türkiye

² Constructive Mathematical Analysis Research Laboratory, Selçuk University, Selçuklu, Türkiye

³ Tuna Edutech Software Ltd., Konya Teknokent, Selçuklu, Türkiye

metinturgay@yahoo.com

key-words: Bivariate sampling operators, simultaneous approximation, weighted spaces.

Abstract:

This study introduces a theoretical framework for bivariate Kantorovich-type sampling operators in weighted function spaces. We investigate the rate of simultaneous approximation using multidimensional Taylor expansions and mixed moduli of smoothness. We demonstrate that the mathematical derivative of the operator smoothly approaches the generalized derivative of the function. Furthermore, we discuss the practical implications of this theoretical property in digital image processing. Since digital images are discrete structures that are inherently non-differentiable, classical edge detection methods often amplify noise. We present a methodological concept where the derivative of the continuous operator kernel serves as an analytical edge detector. This mathematical architecture proposes a deterministic, one-step method for simultaneous image denoising and feature extraction, bridging pure approximation theory with potential applications in medical imaging analysis.

References

- [1] Acar, T., Costarelli, D. and Vinti, G. (2020). Linear prediction and simultaneous approximation by m -th order Kantorovich type sampling series. *Banach J. Math. Anal.*, 14, 1481–1508.
- [2] Bardaro, C., Vinti, G., Butzer, P.L. et al. (2007). Kantorovich-Type Generalized Sampling Series in the Setting of Orlicz Spaces. *STSIP*, 6, 29–52.
- [3] Draganov, B. R., Weighted Simultaneous Approximation by Kantorovich Sampling Operators. In: *Proc. intern. conf. "Constructive Theory of Functions", Lozenets 2023*, B. Draganov, K. Ivanov, G. Nikolov, R. Uluchev (eds.), Prof. Marin Drinov Academic Publishing House, Sofia, 2024, 17-41.
- [4] Rinelli, L., Travaglini, A., Vescera, N. and Vinti, G. (2026). Sampling type operators versus AI in medical image processing. *Discrete and Continuous Dynamical Systems - S*, 28, 216-242.

4th International Conference: Constructive Mathematical Analysis

Risk-Aware Autonomous Decision Analysis under Cross-Domain Distribution Shift Using Uncertainty and Explanation Entropy

Emine Ceren Karademir¹, Gökhan Uçkan¹, Arife Çınar¹ and Emre Çomak¹

¹ Department of Computer Engineering, Pamukkale University, Denizli, Türkiye

ekarademir22@posta.pau.edu.tr, guckan@pau.edu.tr, acinar211@posta.pau.edu.tr, ecomak@pau.edu.tr

key-words: Deep Ensemble, domain shift, epistemic uncertainty, focus entropy, explainable AI, risk-aware triage, diabetic retinopathy

Abstract:

Deep learning-based diabetic retinopathy (DR) screening may become unreliable under domain shift across acquisition settings and populations [4]. This study proposes an entropy- and uncertainty-based framework for identifying potentially erroneous autonomous decisions under cross-domain shift.

For each input x , let $\mathcal{M} = \{f_m\}_{m=1}^M$ denote an ensemble of M probabilistic classifiers. Ensemble outputs yield epistemic uncertainty, maximum confidence, and prediction margin [2]. Grad-CAM and SmoothGrad maps are generated, and focus entropy measures the dispersion of model attention. These signals are integrated into an XGBoost triage model that estimates whether an autonomous decision is likely to be incorrect.

The framework was evaluated using EyePACS, APTOS 2019, and Messidor-2, with IDRiD used for explanation-based validation. The base system comprised three EfficientNet-B0 classifiers [3]. A model trained on EyePACS [1] achieved 82.39% source-domain accuracy but 69.15% on Messidor-2, with recall decreasing to 40.72%. Incorrect predictions could remain overconfident, and explanation maps frequently shifted from retinal lesions toward less informative regions.

The risk filter achieved 94.75% AUC. Of 3,662 images, 2,485 (68%) were assigned to a low-risk autonomous group with no observed errors; the remaining 1,177 (32%) were referred for specialist review and contained all 512 base-model errors. Ensemble uncertainty, focus entropy, and risk triage therefore provide an interpretable reliability layer for autonomous screening under domain shift.

References

- [1] Cuadros, J., Bresnick, G. (2009). EyePACS: An adaptable telemedicine system for diabetic retinopathy screening. *Journal of Diabetes Science and Technology*, 3(3), 509–516.
- [2] Lakshminarayanan, B., Pritzel, A. and Blundell, C. (2017). Simple and scalable predictive uncertainty estimation using deep ensembles. *Advances in Neural Information Processing Systems*, 30.
- [3] Tan, M., Le, Q. (2019). EfficientNet: Rethinking model scaling for convolutional neural networks. *International Conference on Machine Learning*, pp. 6105–6114.
- [4] Ting, D. S. et al. (2019). Deep learning for diabetic retinopathy screening: A clinical translation perspective. *The Lancet Digital Health*, 1(1), e40–e53.

4th International Conference: Constructive Mathematical Analysis

Approximation by Generalized Baskakov Operators Associated with Exponential Functions

Gülsüm Ulusoy Ada¹, Mussa Yousef Mohamed¹

¹ Institute of Graduate Studies, Çankırı Karatekin University, Çankırı, Türkiye

gulsumulusoy@karatekin.edu.tr, mussablhaj1980@gmail.com

key words: Baskakov operators, Stancu-type extension, exponential functions, uniform convergence, Voronovskaya-type asymptotic theorem.

Abstract:

This study introduces a novel Stancu-type extension for a specific class of Baskakov operators frameworks linked with exponential functions. Initially, the paper delves into the core geometric and algebraic properties of these newly constructed operators, providing precise computations for their moments. By employing exponential test functions, the uniform convergence of these operators within the Banach space $C^*[0, \infty)$ is rigorously demonstrated.

Furthermore, the manuscript evaluates the qualitative degree of approximation, establishing quantitative bounds using the framework of the exponential modulus of continuity. Lastly, a Voronovskaya-type asymptotic theorem is formulated and proved to characterize the first-order asymptotic expansion of the operators. The insights and theorems established in this research offer meaningful advancements to the field of positive linear operators, effectively generalizing previous literature on exponential-preserving Baskakov-type operators used for alignment for the different equations

References

- [1] Baskakov, V. A. (1957). An example of a sequence of linear positive operators in the space of continuous functions. Dokl. Akad. Nauk SSSR, 113, 249–251.
- [2] Boyanov, B. D., Veselinov, V. M. (1970). A note on the approximation of functions in an infinite interval by linear positive operators. Bull. Math. Soc. Sci. Math. Roumanie, 14(62), 9–13.
- [3] Bodur, M., Gurel Yilmaz, O. and Aral, A. (2018). Approximation by Baskakov-Szász-Stancu operators preserving exponential functions. Construct. Math. Anal., 1(1), 1–8.
- [4] Holhos, A. (2010). The rate of approximation of functions in an infinite interval by positive linear operators. Stud. Univ. Babeş-Bolyai Math., 55(2), 133–142.
- [5] Stancu, D. D. (1968). Approximation of functions by a new class of linear polynomial operators. Rev. Roumaine Math. Pures Appl., 13, 1173–1194.

4th International Conference: Constructive Mathematical Analysis

An Inertial Projection-Based Neurodynamic Approach for Image Restoration and Inpainting

Rattanaorn Wangkeeree¹, Rabian Wangkeeree¹, Kasamsuk Ungchittrakool¹
and Yirga Abebe Belay^{1,2}

¹ Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, Thailand

² Department of Mathematics, Aksum University, Axum, Ethiopia

rattanaornw@nu.ac.th, rabianw@nu.ac.th, kasamsuku@nu.ac.th, yirga2006@gmail.com

key-words: Inertial neurodynamic model, projection-based RNN, image inpainting, image restoration, lyapunov stability.

Abstract:

The ℓ_1 -regularized least squares optimization, widely known as the LASSO problem, is a fundamental mathematical framework for sparse signal recovery, image restoration, and image inpainting. While recurrent neural networks (RNNs) provide an effective real-time parallel computing environment for such tasks, existing neurodynamic models often face limitations due to high computational overhead in dimension-doubling variable splitting or slow convergence in first-order architectures. To address these challenges, this paper proposes a novel Inertial Projection-Based Recurrent Neural Network (IP-RNN). By integrating a Prox-Projection identity to manage the non-differentiable ℓ_1 subgradient within the original problem space and incorporating a second-order inertial extrapolation term to accelerate the convergence rate, the proposed model achieves significant computational efficiency. Through a rigorous theoretical analysis using Lyapunov stability theory, we prove the global asymptotic stability of the system, ensuring its convergence to the optimal solution. Numerical experiments on various image restoration and inpainting tasks demonstrate that the proposed IP-RNN consistently achieves competitive results in terms of Signal-to-Noise Ratio (SNR) and Structural Similarity (SSIM) indices while maintaining a more compact and faster architecture compared to conventional neurodynamic approaches.

References

- [1] Arfat, Y., Kumam, P., Khan, M. A. A. and Sa Ngiamsunthorn, P. (2022). Shrinking approximants for fixed point problem and generalized split null point problem in Hilbert spaces. *Optim. Lett.*, 16, 1895-1913.
- [2] Mohammadi, M., Atashin, A. A. and Tamburri, D. A. (2023). From l_1 subgradient to projection: A compact neural network for l_1 -regularized logistic regression. *Neurocomputing*, 526, 30-41.
- [3] Mohammadi, M., Tan, Y.-H., Hofman, W. and Mousavi, S. H. (2018). A novel one-layer recurrent neural network for the l_1 -regularized least square problem. *Neurocomputing*, 315, 135-144.
- [4] Reich, S., Tuyen, T. M. (2020). Parallel iterative methods for solving the generalized split common null point problem in Hilbert spaces. *RACSAM*, 114, 180.
- [5] Yodjai, P., Kumam, P., Kitkuan, D., Jirakitpuwapat, W. and Plubtieng, S. (2019). The Halpern approximation of three operators splitting method for convex minimization problems with an application to image inpainting. *Bangmod Int. J. Math. Comp. Sci.*, 5(1-2), 58-75.

4th International Conference: Constructive Mathematical Analysis

Weinstein Pseudo-Differential Operators on Torus

Santosh Kumar Upadhyay

Department of Mathematical Sciences, Indian Institute of Technology (BHU) Varanasi, Varanasi, India

skupadhyay.apm@itbhu.ac.in

key-words: Pseudo-differential operators, Weinstein transform, discrete Weinstein transform.

Abstract:

Pseudo-differential operators are combinations of integral transform and partial differential operators on $\mathbb{T}^{n+1}$. This theory plays a significant role in various problems involving partial differential operators by exploiting the discrete Weinstein transform on $\mathbb{Z}^{n+1}$. In this talk, the concept and various properties of the Weinstein pseudo-differential operators will be discussed in relation to the discrete Weinstein transform.

References

- [1] Botchway, L.N.A., Kibiti, P.G. and Ruzhansky, M. (2020). Difference equations and pseudo-differential operators on $\mathbb{Z}^n$. J. Funct. Anal., 278(11), 108473.
- [2] Ruzhansky, M., Turunen, V. (2010). Quantization of pseudo-differential operators on the torus. J. Fourier Anal. Appl., 16(6), 943–982.
- [3] Upadhyay, S.K., Sartaj, M. (2022). An integral representation of pseudo-differential operators involving the Weinstein transform. J. Pseudo-Differ. Oper. Appl., 13(1), 11.
- [4] Wong, M.W. (2014). An Introduction to Pseudo-Differential Operators. 3rd edn. World Scientific Publishing Company, Singapore.

4th International Conference: Constructive Mathematical Analysis

The Difference of Squares Formula and Quadratic Equations

Syrgak Kydyraliev¹, Anarkul Urdaletova², Elena Burova¹ and Saltanat Japarova³

¹ Department of Applied Mathematics and Informatics, American University of Central Asia, Bishkek, Kyrgyz Republic

² Department of Mathematics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyz Republic

³ Department of Mathematics, Issyk-Kul State University named after Kasym Tynystanov, Karakol, Kyrgyz Republic.

kydyraliev_s@auca.kg, anarkul.urdaletova@manas.edu.kg, burova_e@auca.kg, japarova@iksu.kg

key-words: Difference of squares, quadratic equations, alternative solving methods.

Abstract:

The history of solving quadratic equations stretches back over four thousand years to Ancient Babylon, where scribes calculated land areas and astronomical cycles using rules remarkably similar to our own. However, most cuneiform texts provide only "recipes"—step-by-step instructions without explaining the underlying logic. Paradoxically, modern mathematics education often mirrors this ancient approach, favoring universal formulas (the "standard recipe") while neglecting more elegant and efficient alternatives. This study demonstrates that a wide range of problems typically solved via standard quadratic equations can be addressed more effectively through the direct application of the difference of squares formula.

We propose a specific algorithm based on this approach, highlighting its relevance for mathematical modeling. Many real-world problems, when initially translated into mathematical language, naturally take a form that is ideally suited for the difference of squares method before they are forcibly reduced to a standard quadratic form. By revisiting techniques once championed by Diophantus, Al-Khwarizmi, and Fibonacci—methods largely overshadowed since the 17th century—this work offers a streamlined alternative to the "general recipe," enhancing both computational efficiency and conceptual clarity in the classroom.

References

- [1] Kydyraliev, S. K., Urdaletova, A., Japarova, S. N. and Burova, E. (2025). From Diophantus and Al-Khwarizmi to the Modern Times. Tech Fusion in Business and Society. Studies in Systems, Decision and Control, 233, 335–339.
- [2] Kydyraliev, S. K., Urdaletova, A. B. (2017). Living Mathematics. 7th grade. Bishkek: Solution.
- [3] Kydyraliev, S. K., Urdaletova, A. B. (2020). Fashionable formula. Vestnik KRSU, 20(4), 3–9.

4th International Conference: Constructive Mathematical Analysis

Multi-State Performance Analysis of Benes Network

S. Vishal Vignesh¹, Gundala Vijayalakshmi¹

¹ Department of Mathematics, College of Engineering and Technology, SRM Institute of Science and Technology, Kattankulathur, Tamil Nadu, India

vs8143@srmist.edu.in, vijayalg@srmist.edu.in

key-words: Multi-state Benes network, reliability, availability.

Abstract:

The multi-state Benes network is a widely used interconnection architecture in communication and parallel processing systems due to its fault-tolerant and rearrangeable non-blocking characteristics. This paper presents a comprehensive reliability and performance analysis of a multi-state Benes network using a combined Markov Universal Generating Function (CMUGF) approach. The proposed methodology integrates continuous-time Markov processes with the UGF technique to effectively model the stochastic behavior and multiple performance states of switching elements and links under failure and repair mechanisms. A generalized CMUGF algorithm is developed to evaluate important system performance measures, such as reliability, availability, mean time to failure (MTTF), sensitivity analysis and Birnbaum importance measures. The proposed algorithm successfully computes all targeted performance measures for the constant failure and repair rates while efficiently handling the large state-space complexity associated with multi-state systems. Numerical results are compared with existing research work to validate the accuracy and effectiveness of the proposed methodology. The comparative analysis demonstrates improved computational efficiency and enhanced reliability assessment capability of the developed framework.

References

- [1] Bistouni, F., Jahanshahi, M. (2014). Improved extra group network: a new fault-tolerant multistage interconnection network. *J Supercomput*, 69, 161–199.
- [2] Blake, J. T., Trivedi, K. S. (1988). Multistage interconnection network reliability. *IEEE Transactions on Computers*, 38(11), 1600–1604.
- [3] Jahanshahi, M., Bistouni, F. (2015). Improving the reliability of the Benes network for use in large-scale systems. *Microelectronics Reliability*, 55(3), 679–695. 10.1016/j.microrel.2014.12.008.
- [4] Khati, A., Bisht, S. (2025). Chapter 15 - Reliability analysis of interconnection networks: A comprehensive review of Shuffle-exchange, Benes, and Gamma networks. In A. Kumar, A. S. Bhandari and M. Ram (Eds.), *Reliability Assessment and Optimization of Complex Systems* (pp. 287–308). Elsevier. 10.1016/B978-0-443-29112-8.00016-5
- [5] Wu, C. L., Feng, T. Y. (1980). On a class of multistage interconnection networks. *IEEE Transactions on Computers*, 29(8), 694–702.

4th International Conference: Constructive Mathematical Analysis

Convergence of the Generalised Picard-Mann Iteration for G-Monotone Non-Expansive Mappings in Banach Spaces with a Graph

Deepa Vijayaseenan¹, Rekha Panicker¹ and Rahul Shukla¹

¹ Department of Mathematical Sciences and Computing, Yunivesithy Walter Sisulu, Mthatha, South Africa

223056162@mywsu.ac.za, rpanicker@wsu.ac.za, rshukla@wsu.ac.za

Key-words: Generalised Picard-Mann iteration, G-monotone nonexpansive mappings, uniformly convex Banach space.

Abstract:

In this paper, we study the convergence of the generalised Picard-Mann iteration for approximating fixed points of G -monotone nonexpansive mappings in uniformly convex Banach spaces endowed with a directed graph structure. A mapping $T : C \rightarrow C$ on a nonempty closed convex subset C of a Banach space X is called G -monotone nonexpansive if it preserves the edge relation of a directed graph $G = (V(G), E(G))$ with $V(G) = C$ and satisfies $\|Tx - Ty\| \leq \|x - y\|$ for all $(x, y) \in E(G)$. The iterative scheme under investigation is defined by $x_{n+1} = T^k((1 - \alpha_n)x_n + \alpha_n Tx_n)$, where $k \geq 1$ is a fixed positive integer and $\{\alpha_n\} \subset (0, 1)$ is a control sequence. We first observe that the iteration developed by Wen (2019) is recovered as a special case when $k = 2$, thereby establishing that the generalised Picard-Mann scheme is strictly more general. We then prove two edge propagation lemmas which rigorously verify that the graph structure, specifically the membership of the iterates and their images in the edge set $E(G)$, is preserved throughout the iteration under the assumptions that G is connected and transitive and that G -intervals are convex. These lemmas are essential for the valid application of the G -monotone nonexpansive condition at each step and address a gap that is often treated only informally in the existing literature. Using these results, we establish three main convergence theorems. First, we prove a necessary and sufficient condition: $F(T) \neq \emptyset$ with $(x_0, p) \in E(G)$ for some $p \in F(T)$ if and only if the iterative sequence $\{x_n\}$ is bounded and satisfies $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$, provided $\lim_{n \rightarrow \infty} \alpha_n(1 - \alpha_n) > 0$. Second, we prove weak convergence of $\{x_n\}$ to a fixed point under Opial's property and Property (P). Third, we establish strong convergence when the mapping is additionally semicompact, including a proof that $F(T)$ is closed. Finally, we present a numerical example and demonstrate through comparative computations that the generalised Picard-Mann iteration converges significantly faster than the Wen iteration to the fixed point.

References

- [1] Shukla, R., Pant, R. and Sinkala, W. (2022). A general Picard-Mann iterative method for approximating fixed points of nonexpansive mappings with applications. *Symmetry*, 14, 1741.
- [2] Wen, D.J. (2019). Weak and strong convergence theorems of G -monotone nonexpansive mapping in Banach spaces with a graph. *Numer. Funct. Anal. Optim.*, 40, 163-177.

4th International Conference: Constructive Mathematical Analysis

A New Gradient-Based Neurodynamic Network for Circular Cone Constrained Variational Inequality Problems

Rattanaorn Wangkeeree¹, Rabian Wangkeeree¹, Yirga Abebe Belay², Kasamsuk Ungchittrakool¹ and Nutt Tananimit³

¹ Department of Mathematics, Naresuan University, Phitsanulok, Thailand

² Department of Mathematics, Aksum University, Axum, Ethiopia

³ Department of Mathematics Education, Walailak University, Nakhon Si Thammarat, Thailand

rattanaornw@nu.ac.th, rabianw@nu.ac.th, yirga2006@gmail.com, kasamsuku@nu.ac.th

key-words: Circular cones, neurodynamic networks, variational inequalities.

Abstract:

In this study, we propose a new gradient-based neurodynamic network for solving variational inequality problems constrained by circular cones. Circular cones generalize second-order cones by allowing adjustable aperture parameters, thereby providing greater modeling flexibility in constrained optimization problems. By reformulating the circular cone constrained variational inequality problem into an equivalent system of smoothed equations, we design a steepest-descent neurodynamic system whose equilibrium points correspond exactly to the solutions of the original problem. Using Lyapunov stability theory, we establish the global stability and convergence of the proposed network. Numerical experiments are presented to demonstrate the effectiveness and feasibility of the proposed approach.

References

- [1] Nazemi, A., Sabeghi, A. (2019). A novel gradient-based neural network for solving convex second-order cone constrained variational inequality problems. *J. Comput. Appl. Math.* 347, 343–356.
- [2] Tang, J., Zhou, J. (2020). Smoothing inexact Newton method based on a new derivative-free non-monotone line search for the NCP over circular cones. *Ann. Oper. Res.* 295(2), 787–808.

4th International Conference: Constructive Mathematical Analysis

A Projection Recurrent Neural Network for Circular Cone Constrained Variational Inequalities

Rattanaporn Wangkeeree¹, Rabian Wangkeeree¹, Alireza Nazemi², Yirga Abebe Belay³ and Kasamsuk Ungchitrakool¹

¹ Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

² Department of Mathematics, School of Mathematical Sciences, Shahrood University of Technology, Shahrood, Iran

³ Department of Mathematics, Aksum University, Axum, Ethiopia

rattanapornw@nu.ac.th, rabianw@nu.ac.th, nazemi20042003@gmail.com,
yirga2006@gmail.com, kasamsuku@nu.ac.th

key-words: Circular cone, variational inequality, projection recurrent neural network.

Abstract:

In this paper, we develop a projection recurrent neural network for solving nonlinear variational inequality problems constrained by circular cones, together with related optimization models. Since the circular cone extends the second-order cone, it offers a more general framework for the formulation and analysis of constrained problems. By employing a smoothing technique, the original variational inequality problem is reformulated as an equivalent circular cone programming problem. Based on the relationship between optimal solutions of circular cone programming and the associated Karush–Kuhn–Tucker conditions, we construct a projection-based recurrent neural network with reduced dimensionality and lower computational complexity. Moreover, Lyapunov stability theory is applied to show that the proposed neural network is Lyapunov stable and globally convergent to a solution satisfying the reduced optimality conditions. Numerical examples are included to demonstrate the effectiveness and practical applicability of the proposed approach.

References

- [1] Nazemi, A., Sabeghi, A. (2019). A novel gradient-based neural network for solving convex second-order cone constrained variational inequality problems. *J. Comput. Appl. Math.*, 347, 343–356.
- [2] Tang, J., Zhou, J. (2020). Smoothing inexact Newton method based on a new derivative-free non-monotone line search for the NCP over circular cones. *Ann. Oper. Res.*, 295(2), 787–808.

4th International Conference: Constructive Mathematical Analysis

The Numerical Range of the Generalized Inverse and Its Applications to Iterative Methods

Gerald Wanjala

Department of Mathematics, Sultan Qaboos University/Muscat, Oman

gwanjala@squ.edu.om

key-words: Numerical range, generalized inverse, EP operators, steepest descent.

Abstract:

The numerical range (field of values) of a bounded linear operator is a convex set in the complex plane that contains the spectrum—a property known as spectral inclusion. For singular or ill-conditioned operators, however, the standard spectral inclusion can be exceedingly loose, and traditional inverse theory fails. This talk investigates how the *generalized inverse*, defined variationally as the unique least squares solution of minimal norm [2], can be used to sharpen spectral inclusion and to analyse iterative methods for singular linear equations.

In this talk, we establish a geometric characterization of the closed range condition (which is an essential hypothesis for a bounded generalized inverse) via a gap at zero in the numerical range of the associated positive operator $\bar{T} = T^*T|_{\mathcal{R}(T^*)}$. Secondly, for EP operators (those satisfying $TT^\dagger = T^\dagger T$), we prove a sharpened spectral inclusion:

$$\sigma(T) \subseteq \overline{W(T)} \cap \frac{1}{\overline{W(T^\dagger)}},$$

and we characterize EP operators geometrically by the reciprocity of the numerical ranges of T and $T^\dagger$ on the reducing subspace $\mathcal{R}(T^*)$.

Finally, we show that the convergence of the Petryshyn, Schulz, steepest descent, and conjugate gradient methods for computing $T^\dagger b$, is governed entirely by the numerical range of $\bar{T}$. The convergence factors for these methods, including the optimal parameter for the Petryshyn iteration and the error reduction factor for steepest descent are expressed directly in terms of the bounds

$$m_W = \inf \operatorname{Re}(W(\bar{T})) = \|T^\dagger\|^{-2}, \text{ and } M_W = \sup \operatorname{Re}(W(\bar{T})) = \|T\|^2.$$

References

- [1] Chien, M. T. (2020). Numerical range of Moore-Penrose inverse matrices. *Mathematics*, 8(5), 830.
- [2] Groetsch, C. W. (1977). *Generalized Inverses of Linear Operators: Representation and Approximation*. Marcel Dekker, New York.
- [3] Gustafson, K. E., Rao, D. K. M. (1997). *Numerical Range: The Field of Values of Linear Operators and Matrices*. Springer, New York.

4th International Conference: Constructive Mathematical Analysis

A New Approach to Decision Making Using Soft Covering Based Rough Sets by Concept of Soft Maximal Description

Zehra Güzel Ergül¹, Kubilay Yaşar¹

¹ Departments of Mathematics, Kırşehir Ahi Evran University, Kırşehir, Türkiye

zguzel@ahievran.edu.tr, yasar.kubilay@ahievran.edu.tr

key-words: Soft rough set, soft covering based rough set, decision making.

Abstract:

Soft covering based rough set theory has become a useful and well-known area of research in theories of uncertainty. The present work follows up on this flourishing research topic. While soft covering based rough sets defined via the concept of soft minimal description have been extensively used by various authors in the literature for decision making problems, soft covering based rough sets defined via the concept of soft maximal description have not yet been applied in this context. Motivated by this gap, the present study applies these models within different decision making frameworks. In all cases, the best alternative is selected from a finite set, and in some methods, a ranking of alternatives is also produced. Also, the obtained results are compared in order to evaluate the effectiveness of the considered decision making approaches.

References

- [1] Atagun, A. O., Kamacı, H. and Oktay, O. (2018). Reduced soft matrices and generalized products with applications in decision making. *Neural Comput. and Applic.*, 29, 445-456.
- [2] Atagun, A. O., Kamacı, H. (2024). Decompositions of soft sets and soft matrices with applications in group decision making. *Scientia Iranica*, 31(6), 518-534.
- [3] Cagman, N., Enginoglu, S. (2010). Soft matrix theory and its decision making. *Computer and Mathematics with Applications*, 59, 3308-3314.
- [4] Feng, F., Changxing, L., Davvaz, B. and Ali, M. I. (2010). Soft sets combined with fuzzy sets and rough sets: A tentative approach. *Soft Comput.*, 14, 899-911.
- [5] Feng, F. (2011). Soft rough sets applied to multicriteria group decision making. *Ann. Fuzzy Math. Inform.*, 2(1), 69-80.
- [6] Guzel Ergul, Z., Yuksel, S. (2019). A new type of soft covering based rough sets applied to multicriteria group decision making for medical diagnosis. *Math. Sci. and Appl. E-Notes*, 7(1), 28-38.
- [7] Guzel Ergul, Z., Yasar, K. (2025). New kinds of soft covering based rough sets via the concept of soft maximal description. *Advanced Studies:Euro-Tbilisi Mathematical Journal*, Accepted, In press.
- [8] Molodtsov, D. (1999). Soft set theory-first results. *Comput. Math. Appl.*, 37, 19-31.
- [9] Pawlak, Z. (1982). Rough sets. *International Journal of Computer and Information Sciences*, 11(5), 341-356.
- [10] Yuksel, S, Tozlu, N., and Dizman, T. (2015). An application of multicriteria group decision making by soft covering based rough sets. *Filomat*, 29, 209-219.

- [11] Yuksel S, Guzel Ergul, Z. and Tozlu, N. (2014). Soft covering based rough sets and their application. The Scientific World Journal, Article ID 970893, 9 pages.
- [12] Zhan, J., Alcantud, J. C. R. (2019). A novel type of soft rough covering and its application to multicriteria group decision making. Artificial Intelligence Review, 52, 2381-2410.

4th International Conference: Constructive Mathematical Analysis

Hyperbolic Pythagorean Hodograph Curves

Aziz Yazla¹, Fatma Nur Danis²

¹ Department of Mathematics, Selcuk University, Konya, Türkiye

² Institute of Science, Selcuk University, Konya, Türkiye

azizyazla@gmail.com, fatmanurdanis01@gmail.com

key-words: Pythagorean hodograph curve, Poincaré upper half-plane, Poincaré metric.

Abstract:

In this paper we introduce and completely classify hyperbolic Pythagorean hodograph (H-PH) curves in the Poincaré upper half-plane. Unlike the Euclidean case, where polynomial PH curves form a rich closed class, we show that rational functions constitute the natural and maximal class in the hyperbolic setting. We prove a necessary and sufficient condition for the existence of rational H-PH curves and derive a canonical normal form. Besides, we demonstrate the existence of unit speed rational H-PH curves as a property distinct from the Euclidean case. A precise comparison with Euclidean PH curves is also established.

References

- [1] Anderson, J. W. (2005). Hyperbolic Geometry. Springer, USA.
- [2] Farouki, R.T. (2008). Pythagorean-Hodograph Curves. Springer, Berlin, Heidelberg.
- [3] Farouki, R.T., Sakkalis, T. (1990). Pythagorean Hodographs. IBM Journal of Research and Development, 34(5), 736-752.
- [4] Farouki, R. T., Sakkalis, T. (1991). Real Rational Curves are not “unit speed”. Computer Aided Geometric Design, 8(2), 151-157.
- [5] Stahl, S. (1993). The Poincaré Half-Plane: A Gateway to Modern Geometry. Jones and Bartlett Learning.
- [6] Yazla, A. (2023). Characterization of Minkowski Pythagorean Hodograph Curves with Rational Rotation Minimizing Frames, Ph.D Thesis, Selçuk University.

4th International Conference: Constructive Mathematical Analysis

Characterization of Some Classes of Compact Operators on Cesàro Difference Series Spaces

Buse Yumruktay¹, Canan Hazar Güleç¹

¹ Mathematics, Pamukkale University, Denizli, Türkiye

buse.tercan.bt@gmail.com, gchazar@pau.edu.tr

key-words: Cesàro means, difference operator, series spaces, compact operators.

Abstract:

In this study, it is aimed to compute the norms and the Hausdorff measures of noncompactness of matrix operators acting on the Cesàro difference series space $|C_\alpha|_p(\nabla)$, which is generated by means of the difference operator and the Cesàro mean [1]. Furthermore, by applying the Hausdorff measure of noncompactness, we characterize certain classes of compact operators on Cesàro difference series space. Finally, by making use of these results, some known classical results in the literature are generalized.

References

- [1] Güleç, G. C. H. (2020). Difference series spaces and matrix transformations. Türk Doğa ve Fen Dergisi, 9(1), 137-143.
- [2] Hazar, G. C., Sarıgöl M. A. (2018). Absolute Cesàro series spaces and matrix operators. Acta App. Math., 154, 153-165.
- [3] Hazar Güleç, G. C. (2020). Compact matrix operators on absolute Cesàro spaces. Numerical Functional Analysis and Optimization, 41(1), 1-15.

This study was supported by the Pamukkale University Scientific Research Projects Unit with project number 2026FEBE005.

4th International Conference: Constructive Mathematical Analysis

Modules with ss -Supplements in Their Cofinite Extensions

Başak Zaman¹, Emine Önal Kır¹

¹ Department of Mathematics, Kırşehir Ahi Evran University, Kırşehir, Türkiye

basakzzaman@gmail.com, emine.onal@ahievran.edu.tr

key-words: Semisimple modules, ss -supplement submodules, cofinite extensions, semiperfect rings, CE_{ss} -supplementing modules.

Abstract:

The concept of CE_{ss} -supplementing modules is introduced and analyzed. A module T is termed CE_{ss} -supplementing whenever each cofinite extension of T contains an ss -supplement of T . It is demonstrated that for modules over commutative von Neumann regular rings, the properties of cofinite injectivity, being CE_{ss} -supplementing, and possessing property (CE) coincide. Consequently, for left V -rings, it follows that a module is CE_{ss} -supplementing precisely when it has property (CE) . The property CE_{ss} -supplementing transfers to direct summands. The amply CE_{ss} -supplementing property of a module is equivalent to the requirement that every submodule inherits the CE_{ss} -supplementing condition. It is proved that a ring S is semiperfect with $Rad(_S S) \leq Soc(_S S)$ if and only if every S -module is (amply) CE_{ss} -supplementing.

References

- [1] Çalışıcı, H., Türkmen, E. (2012). Modules that have a supplement in every cofinite extension. Georgian Mathematical Journal, 19, 209-216.
- [2] Kaynar, E., Çalışıcı, H. and Türkmen, E. (2020). ss -Supplemented modules. Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, 69(1), 473-485.
- [3] Sharpe, D. W., Vamos, P. (1972). Injective Modules. Lecturers in Pure Mathematics University of Sheffield, Cambridge at the University Press.



<https://iccma2026.selcuk.edu.tr/>
<https://dergipark.org.tr/en/pub/cma>

CMA

CONSTRUCTIVE MATHEMATICAL ANALYSIS